

EXACYL®

Tranexamic acid

EXACYL film-coated tablets:

- Box of 20 tablets each containing 250 mg.
- Box of 20 tablets each containing 500 mg.

EXACYL oral solution :

- Box of 5 ampules of 10 ml each containing 1 g.

INDICATIONS:

This drug is an ANTIHAEMORRHAGICAL agent recommended for bleedings.

CONTRAINDICATIONS:

This drug is contraindicated in patients with:

- Thombosis,
- Severe renal insufficiency.

In case of doubt, consult your doctor or pharmacist.

PRECAUTIONS :

Inform your doctor in the event of pregnancy or breast-feeding.

NEVER LEAVE THE DRUG WITHIN THE REACH OF CHILDREN.

OTHER POSSIBLE SIDE EFFECTS OF THIS DRUG:

As with any active product, this drug can, in some individuals, produce more or less troublesome side effects: nausea, vomiting, vertigo, etc.

DOSAGE AND ADMINISTRATION:

ORAL ROUTE

EXACYL film-coated tablets:

- Adults : 2 to 4 g per 24 hours in 2 or 3 divided doses.
- Children : 20 mg/kg per day on average.

EXACYL oral solution:

- Adults : 2 to 4 ampules per 24 hours, divided in several doses.
- Children : follow strictly your doctor's prescription.

IN ALL CASES FOLLOW STRICTLY YOUR DOCTOR'S PRESCRIPTION.

DISPENSING CONDITION:

This drug is registered on list I

Your pharmacist can deliver it again only if it has been specified on the prescription, or on a new prescription.

This drug has been prescribed for you personally, in a specific situation.

- It may not be appropriate for another case.
- Do not reuse it without medical advice.
- Do not recommend it to another person.

SHELF-LIFE:

Do not use after expiry date indicated on the packaging.