

**Denk  
Pharma**

# DOUZABIN RETARD FORTE

## Instructions for use

Please read carefully

### Injection

Ampoules for parenteral use.

Each ampoule of 2 ml contains;  
Hydroxocobalamin 10.000 mcg

### Indications

Algias of different origin; algic multiple sclerosis; metabolic, toxic or infectious polyneuritis, zonos and their painful complications, hyperalgic sciatica, cervico-brachial or crural neuralgia, paraesthetic meralgia, scapulohumeral peri-arthritis, arthritic pains, pains of different and indefinite origin, severe alterations of the general status with proteinic deficiency.

### Contra-indications

Sensitivity to hydroxocobalamin.

As with all medicines careful consideration should be given before the administration of DOUZABIN RETARD FORTE during the first trimester of pregnancy. Vitamins are excreted in the breast milk.

### Adverse effects

Allergic hypersensitivity reactions have occurred rarely following the administration of DOUZABIN RETARD FORTE.

Other adverse effects may include in individual cases, acne.

### Interactions with other medications

Serum concentrations may be decreased by the concurrent administration of oral contraceptives.

## Dosage and administration

Unless otherwise prescribed by the physician one ampoule daily or every second day by **slow intramuscular injection** in series of 6-12 injections.

The duration of treatment should be determined by the physician.

### Storage

Store below 25°C.

Do not use this product after the expiry date.

**Keep out of reach of children.**

E. DENK OHG, 81675 MUENCHEN  
Prinzregentenstrasse 79  
Germany

Vers. 2687/10.2003

421DOUZABIN/1