

Salinal[®]

Chewable Tablets, Drops

Antiflatulent

Tablets - Each tablet contains Simethicone 42 mg.

Drops - Each 1 mL contains Simethicone 41.2 mg.

Properties :

Salinal is highly effective in relieving the discomfort and pain due to gaseous distention in the gastro-intestinal tract. Its active component, Simethicone, is a surface active agent. It acts as a dispersant of gas bubbles by reducing surface tension, destroying thereby mucosal pockets of gas which cause distress.

Indications :

Salinal is indicated to alleviate the painful symptoms of excess gas in the digestive tract. **Salinal** is also a useful adjunct in the treatment of many conditions in which the retention of gas may be a problem, including postoperative gaseous distention. The antifoaming property of **Salinal** can be utilized during gastroscopy and bowel radiography to reduce gas shadow and enhance visualization.

Salinal Drops is mainly indicated for infants and babies to relieve bloating and flatulence resulting from feeding problems, but it is not recommended for infant colics. Adults having difficulties in taking tablets may prefer to use the drops form.

Dosage :

Tablets - 1-2 tablets to be chewed four times a day, after meals and at bedtime.

Drops - Adults and children: 1-2 mL four times a day after meals and at bedtime.

Infants (under 2 years) : 0.5 mL four times daily with

meals and at bedtime, or as directed by the physician. The dosage can be mixed with infant food or other suitable liquid to ease administration.

For diagnostic purposes, 2 chewable tablets or 2 mL drops to be taken 3 times, one day before the examination, and 2 tablets or 2 mL drops in the morning of the day of examination.

Side Effects:

No side effects to this product have been reported.

Presentation:

Tablets : 30 tablets in blisters.

Drops : 15 mL in a bottle with a graduated dropper.

THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep all medicines out of the reach of children.



Produced by: **juphar**
Gulf Pharmaceutical Industries,
Ras Al Khaimah, U. A. E.