

DIFRAREL® E

coated tablet

QUALITATIVE AND QUANTITATIVE COMPOSITION

ANTHOCYANOSIDIC EXTRACT OF BILBERRY	50.000 mg
ALPHATOCOPHEROL ACETATE (concentrate of) - pulverulent form	50.000 mg
Excipient : Gelatine, Silicic Acid, Microcrystalline Cellulose, Sodid Croscarmellose, Anhydrous Colloidal Silica, Arabic Gum, Saccharose, Corn Starch, Talc, Titanium dioxide (E 171), Opaglos 6000.	
For one coated tablet	

PHARMACEUTICAL PRESENTATION

Box of 24 coated tablets.

PHARMACO-THERAPEUTIC CLASSIFICATION

OPHTHALMOLOGICAL MEDICINE / VASCULAR PROTECTANT

(S: sensory organs)

NAME AND ADDRESS OF THE PRODUCER *Laboratoires LEURQUIN MEDIOLANUM*

68-88 rue Ampère - Z.I. des Chanoux

93330 NEUILLY SUR MARNE - FRANCE

Medical information : ☎ 33 (0)1 49 44 29 29

CASE(S) IN WHICH THIS MEDICINE CAN BE USED

This medicine is recommended as a back-up treatment for dusk and night vision disorders, short-sightedness.

WARNING!

CASE(S) IN WHICH THIS MEDICINE SHOULD NOT BE USED

This medicine SHOULD NOT BE USED in the following case:

- previous history of allergy to one of the constituents of the coated tablet.

IF IN DOUBT, DO NOT HESITATE TO CONSULT YOUR DOCTOR OR CHEMIST.

PRECAUTIONS

You should consult your doctor in the event of sudden vision difficulties or worsening of a pre-existent vision problem.

IF IN DOUBT, DO NOT HESITATE TO CONSULT YOUR DOCTOR OR CHEMIST

INTERACTION WITH OTHER MEDICINES AND OTHER INTERACTIONS

TO AVOID POSSIBLE INTERACTION BETWEEN SEVERAL MEDICINES, YOUR DOCTOR OR CHEMIST MUST BE SYSTEMATICALLY INFORMED OF ANY OTHER CURRENT TREATMENT.

PREGNANCY - BREAST FEEDING

It is not advisable to take this medicine during pregnancy.

If you should find that you are pregnant while taking the medicine, consult your doctor who can best decide on the necessity to continue.

Avoid taking this medicine when breast feeding.

Generally speaking, during pregnancy and while breast feeding, you should always consult your doctor or chemist before taking any medicine.

ADMINISTRATION OF THIS MEDICINE

DOSE

3 to 6 tablets a day.

The treatment is to be taken for 20 days per month, then renewed if necessary.

MODE AND METHOD OF ADMINISTRATION

To be taken orally.

Swallow the tablets, without chewing, with a glass of water.

LENGTH OF TREATMENT

The course of treatment should be taken for 20 days per month and renewed if necessary.

Follow your doctor's recommendations.

UNDESIRABLE, TROUBLESOME SIDE-EFFECTS

AS IS THE CASE FOR ANY ACTIVE SUBSTANCES, THIS MEDICINE MAY CAUSE UNDESIRABLE OR TROUBLESOME SIDE-EFFECTS IN CERTAIN INDIVIDUALS;

INFORM YOUR DOCTOR OR CHEMIST OF ANY UNDESIRABLE OR TROUBLESOME SIDE-EFFECT NOT MENTIONED ON THIS LEAFLET.

STORAGE

DO NOT EXCEED THE EXPIRY DATE SHOWN ON THE OUTER PACKAGING.

SPECIAL STORAGE PRECAUTIONS

To store at a temperature not exceeding 30°C.

LEAFLET REVISION DATE : February 2006, 28th