

Reparil®-Dragees



A selective anti-exudative and anti-inflammatory agent.

Composition:

1 dragee contains 20 mg amorphous aescin.

The action of Reparil-Dragees:

Aescin, the active principle of horse chestnut in pure form, has anti-exudative, anti-inflammatory and anti-oedema effects at the site of the lesion. It quickly relieves symptoms caused by localised swelling (sensations of pressure, pain, etc.) without adverse effects on mineral metabolism or the blood-forming system.

When to take Reparil-Dragees:

Varicosis; leg ulcers; haemorrhoids; thrombophlebitis; traumatic swellings (injuries); post-concussional headache; tenosynovitis; painful lesions of the vertebral column. For the continuation of treatment begun with intravenous Reparil.

How to take Reparil-Dragees:

Treatment should be commenced with 2 dragees 3 times daily, taken with fluid after meals. In milder cases and as a maintenance dose 1 dragee should be taken 3 times daily.

Dosage for children:

1 dragee 2 - 3 times daily taken with fluid after meals.

Contra-indications:

Pre-existing renal damage; renal failure; renal insufficiency; Rhesus incompatibility in pregnancy.

Note:

Although animal experiments with aescin have shown no evidence of teratogenic action, in conformity with general medical considerations Reparil should not be given during the first three months of pregnancy.

Presentation:

Prescription packs of 40 and 100 dragees

THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

KEEP MEDICAMENT OUT OF REACH OF CHILDREN

Council of Arab Health Ministers
Union of Arab Pharmacists

(ان هذا الدواء)

- الدواء متغير يؤثر على صحتك واشتراك علاقا للعمليات بمرتكب للملار
- اذبح بدلة وصفة الطبيب وطريقة استعمال المتصور عليها وتعليمات
- الصيدلاني الذي صرفها لك
- فالطبيب والصيدلاني هما الخبراء بالادوية وبخطورتها
- لا تلتصق مدة العلاج الجديدة لك من تلقا نفسك
- لا تكرر صرف الادوية بدون وصفة طبية

لا تدرك الادوية في متناول ايدي الاطفال

مجلس وزراء الصحة العرب
وال اتحاد الصيدلة العرب