

**PACKAGE LEAFLET:  
INFORMATION FOR THE USER**

## Vidaza™ 25 mg/mL powder for suspension for injection

Azacitidine

### Read all of this leaflet carefully before you start using this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, pharmacist or nurse.

#### In this leaflet:

1. What Vidaza is and what it is used for
2. Before you use Vidaza
3. How to use Vidaza
4. Possible side effects
5. How to store Vidaza
6. Further information

## 1. WHAT VIDAZA IS AND WHAT IT IS USED FOR

#### What Vidaza is

Vidaza is an anti-cancer agent which belongs to a group of medicines called 'anti-metabolites'. Vidaza contains the active substance 'azacitidine'.

#### What Vidaza is used for

Vidaza is used in adults who are not able to have a stem cell transplantation to treat:

- higher-risk myelodysplastic syndromes (MDS)
- chronic myelomonocytic leukaemia (CMML)
- acute myeloid leukaemia (AML)

These are diseases which affect the bone marrow and can cause problems with normal blood cell production.

#### How Vidaza works

Vidaza works by preventing cancer cells from growing. Azacitidine becomes incorporated into the genetic material of cells (ribonucleic acid (RNA) and deoxyribonucleic acid (DNA)). It is thought to work by altering the way the cell turns genes on and off and also by interfering with the production of new RNA and DNA. These actions are thought to correct problems with the maturation and growth of young blood cells in the bone marrow that cause myelodysplastic disorders, and to kill cancerous cells in leukaemia.

Talk to your doctor or nurse if you have any questions about how Vidaza works or why this medicine has been prescribed for you.

## 2. BEFORE YOU USE VIDAZA

#### Do not use Vidaza

- if you are allergic to azacitidine or any of the other ingredients of this medicine (listed in section 6).
- if you have advanced liver cancer.
- if you are breast-feeding.

#### Take special care with Vidaza

Talk to your doctor, pharmacist or nurse before using Vidaza in the following situations:

- if you have decreased counts of platelets, red or white blood cells.
- if you have kidney disease.
- if you have liver disease.
- if you have ever had a heart condition or heart attack or any history of lung disease.

#### Blood test

You will have blood tests before you begin treatment with Vidaza and at the start of each period of treatment (called a 'cycle'). This is to check that you have enough blood cells and that your liver and kidneys are working properly.

#### Children and adolescents

Vidaza is not recommended for use in children and adolescents below the age of 18.

### The following information is intended for healthcare professionals only:

#### Recommendations for safe handling

Vidaza is a cytotoxic medicinal product and, as with other potentially toxic compounds, caution should be exercised when handling and preparing azacitidine suspensions. Procedures for proper handling and disposal of anticancer medicinal products should be applied.

If reconstituted azacitidine comes into contact with the skin, immediately and thoroughly wash with soap and water. If it comes into contact with mucous membranes, flush thoroughly with water.

#### Incompatibilities

This medicinal product must not be mixed with other medicinal products except those mentioned below (see "Reconstitution Procedure").

#### Reconstitution procedure

Vidaza should be reconstituted with water for injections. The shelf life of the reconstituted medicinal product can be extended by reconstituting with refrigerated (2 °C to 8 °C) water for injections. Details on storage of the reconstituted product are provided below.

1. The following supplies should be assembled: Vial(s) of azacitidine; vial(s) of water for injections; non-sterile surgical gloves; alcohol wipes; 5 mL injection syringe(s) with needle(s).
2. 4 mL of water for injections should be drawn into the syringe, making sure to purge any air trapped within the syringe.
3. The needle of the syringe containing the 4 mL of water for injections should be inserted through the rubber top of the azacitidine vial followed by injection of the water for injections into the vial.
4. Following removal of the syringe and needle, the vial should be vigorously shaken until a uniform cloudy suspension is achieved. After reconstitution each mL of suspension will contain 25 mg of azacitidine (100 mg/4 mL). The reconstituted product is a homogeneous, cloudy suspension, free of agglomerates. The product should be discarded if it contains large particles or agglomerates. Do not filter the suspension after reconstitution since this could remove the active

#### Other medicines and Vidaza

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. This is because Vidaza may affect the way some other medicines work. Also, some other medicines may affect the way Vidaza works.

#### Pregnancy, breast-feeding and fertility

##### Pregnancy

You should not use Vidaza during pregnancy as it may be harmful to the baby. Use an effective method of contraception during and up to 3 months after treatment. Tell your doctor straight away if you become pregnant during treatment.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

##### Breast-feeding

You should not breast-feed when using Vidaza. It is not known if this medicine passes into human milk.

##### Fertility

Men should not father a child while receiving treatment with Vidaza. Use an effective method of contraception during and up to 3 months after treatment with this medicine.

Talk to your doctor if you wish to conserve your sperm before starting this treatment.

##### Driving and using machines

Do not drive or use any tools or machines if you experience side effects, such as tiredness.

## 3. HOW TO USE VIDAZA

Before giving you Vidaza, your doctor will give you another medicine to prevent nausea and vomiting at the start of each treatment cycle.

- The recommended dose is 75 mg per m<sup>2</sup> body surface area. Your doctor will decide your dose of this medicine, depending on your general condition, height and weight. Your doctor will check your progress and may change your dose if necessary.
- Vidaza is given every day for one week, followed by a rest period of 3 weeks. This "treatment cycle" will be repeated every 4 weeks. You will usually receive at least 6 treatment cycles.

This medicine will be given to you as an injection under the skin (subcutaneously) by a doctor or nurse. It may be given under the skin on your thigh, tummy or upper arm.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## 4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

#### Tell your doctor straight away if you notice any of the following side effects:

- **Drowsiness, shaking, jaundice, abdominal bloating and easy bruising.** These may be symptoms of liver failure and can be life-threatening.
- **Swelling of the legs and feet, back pain, reduced passing of water, increased thirst, rapid pulse, dizziness and nausea, vomiting or reduced appetite and feelings of confusion, restlessness or fatigue.** These may be symptoms of kidney failure and can be life-threatening.
- **A fever.** This could be due to an infection as a result of having low levels of white blood cells, which can be life-threatening.
- **Chest pain or shortness of breath which may be accompanied with a fever.** This may be due to an infection of the lung called "pneumonia", and can be life-threatening.
- **Bleeding.** Such as blood in the stools due to bleeding in the stomach or gut, or such as bleeding inside your head. These may be

symptoms of having low levels of platelets in your blood.

- **Difficulty breathing, swelling of the lips, itching or rash.** This may be due to an allergic (hypersensitivity) reaction.

Other side effects include:

**Very common side effects** (may affect more than 1 in 10 people)

- Reduced red blood count (anaemia). You may feel tired and pale.
- Reduced white blood cell count. This may be accompanied by a fever. You are also more likely to get infections.
- A low blood platelet count (thrombocytopenia). You are more prone to bleeding and bruising.
- Constipation, diarrhoea, nausea, vomiting.
- Pneumonia.
- Chest pain, being short of breath.
- Tiredness (fatigue).
- Injection site reaction including redness, pain or a skin reaction.
- Loss of appetite.
- Joint aches.
- Bruising.
- Rash.
- Red or purple spots under your skin.
- Pain in your belly (abdominal pain).
- Itching.
- Fever.
- Sore nose and throat.
- Dizziness.
- Headache.
- Having trouble sleeping (insomnia).
- Nosebleeds (epistaxis).
- Muscle aches.
- Weakness (asthenia).
- Weight loss.
- Low levels of potassium in your blood.

**Common side effects** (may affect up to 1 in 10 people)

- Bleeding inside your head.
- An infection of the blood caused by bacteria (sepsis). This may be due to low levels of white cells in your blood.
- Bone marrow failure. This can cause low levels of red and white blood cells and platelets.
- A type of anaemia where your red and white blood cells and platelets are reduced.
- An infection in your urine.
- A viral infection causing cold sores (herpes).
- Bleeding gums, bleeding in the stomach or gut, bleeding from around your back passage due to piles (haemorrhoidal haemorrhage), bleeding in your eye, bleeding under your skin, or into your skin (haematoma).
- Blood in your urine.
- Ulcers of your mouth or tongue.
- Changes to your skin at the injection site. These include swelling, a hard lump, bruising, bleeding into your skin (haematoma), rash, itching and changes in the skin colour.
- Redness of your skin.
- Skin infection (cellulitis).
- An infection of the nose and throat, or sore throat.
- Sore or runny nose or sinuses (sinusitis).
- High or low blood pressure (hypertension or hypotension).
- Being short of breath when you move.
- Pain in your throat and voicebox.
- Indigestion.
- Lethargy.
- Feeling generally unwell.
- Anxiety.
- Being confused.
- Hair loss.
- Kidney failure.
- Dehydration.
- White coating covering tongue, inner cheeks, and sometimes on the roof of your mouth, gums and tonsils (oral fungal infection).
- Fainting.
- A fall in blood pressure when standing (orthostatic hypotension) leading to dizziness when moving to a standing or sitting position.
- Sleepiness, drowsiness (somnolence).
- Bleeding due to a catheter line.
- A disease affecting the gut which can result in fever, vomiting and stomach pain (diverticulitis).
- Fluid around the lungs (pleural effusion).
- Shivering (chills).
- Muscle spasms.
- Raised itchy rash on the skin (urticaria).
- Collection of fluid around the heart (pericardial effusion).

**Uncommon side effects** (may affect up to 1 in 100 people)

- Allergic (hypersensitivity) reaction.
- Shaking.
- Liver failure.
- Large plum-coloured, raised painful patches on the skin with fever.
- Painful skin ulceration (pyoderma gangrenosum).

**Rare side effects** (may affect up to 1 in 1,000 people)

- Dry cough.
- Painless swelling in the finger tips (clubbing).
- Tumour lysis syndrome - Metabolic complications that can occur during treatment of cancer and sometimes even without treatment. These complications are caused by the product of dying cancer cells and may include the following: changes to blood chemistry; high potassium, phosphorus, uric acid, and low calcium consequently leading to changes in kidney function, heartbeat, seizures, and sometimes death.

**Not known** (frequency cannot be estimated from the available data)

- Infection of the deeper layers of skin, which spreads quickly, damaging the skin and tissue, which can be life-threatening (necrotizing fasciitis).

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly as described in section 6. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. HOW TO STORE VIDAZA

Keep out of the reach and sight of children. Store below 30 °C

Do not use this medicine after the expiry date which is stated on the vial label and the carton. The expiry date refers to the last day of that month.

Your doctor, pharmacist or nurse are responsible for storing Vidaza. They are also responsible for preparing and disposing of any unused Vidaza correctly.

#### When using immediately

Once the suspension has been prepared it should be administered within 45 minutes.

#### When using later on

If the Vidaza suspension is prepared using water for injections that has not been refrigerated, the suspension must be placed in the refrigerator (2 °C – 8 °C) immediately after it is prepared and kept refrigerated for up to a maximum of 8 hours. If the Vidaza suspension is prepared using water for injections that has been stored in the refrigerator (2 °C – 8 °C), the suspension must be placed in the refrigerator (2 °C – 8 °C) immediately after it is prepared and kept refrigerated for up to a maximum of 22 hours.

The suspension should be allowed up to 30 minutes prior to administration to reach a temperature of approximately (20 °C – 25 °C).

If large particles are present in the suspension it should be discarded.

## 6. FURTHER INFORMATION

#### What Vidaza contains

- The active substance is azacitidine. One vial contains 100 mg azacitidine. After reconstitution with 4 mL of water for injections, the reconstituted suspension contains 25 mg/mL azacitidine.
- The other ingredient is mannitol (E421).

#### What Vidaza looks like and contents of the pack

Vidaza is a white powder for suspension for injection and is supplied in a glass vial containing 100 mg of azacitidine. Each pack contains one vial of Vidaza.

#### Method of administration

Do not filter the suspension after reconstitution.

Reconstituted Vidaza should be injected subcutaneously (insert the needle at a 45-90° angle) using a 25-gauge needle into the upper arm, thigh or abdomen.

Doses greater than 4 mL should be injected into two separate sites.

Injection sites should be rotated. New injections should be given at least 2.5 cm from the previous site and never into areas where the site is tender, bruised, red, or hardened.

Any unused product or waste material should be disposed of in accordance with local requirements.

suspension must be placed in a refrigerator (2 °C to 8 °C) immediately after reconstitution, and kept in the refrigerator for a maximum of 8 hours. If the elapsed time in the refrigerator is greater than 8 hours, the suspension should be discarded appropriately and a new dose prepared.

When reconstituting using refrigerated (2 °C to 8 °C) water for injections, the reconstituted suspension must be placed in a refrigerator (2 °C to 8 °C) immediately after reconstitution, and kept in a refrigerator for a maximum of 22 hours. If the elapsed time in the refrigerator is greater than 22 hours, the suspension should be discarded appropriately and a new dose prepared.

The syringe filled with reconstituted suspension should be allowed up to 30 minutes prior to administration to reach a temperature of approximately 20 °C-25 °C. If the elapsed time is longer than 30 minutes, the suspension should be discarded appropriately and a new dose prepared.

**Calculation of an individual dose**  
The total dose, according to the body surface area (BSA) can be calculated as follows:

Total dose (mg) = Dose (mg/m<sup>2</sup>) x BSA (m<sup>2</sup>)

The following table is provided only as an example of how to calculate individual azacitidine doses based on an average BSA value of 1.8 m<sup>2</sup>.

Dose mg/m <sup>2</sup> (% of recommended starting dose)	Total dose based on BSA value of 1.8 m <sup>2</sup>	Number of vials required	Total volume of reconstituted suspension required
75 mg/m <sup>2</sup> (100 %)	135 mg	2 vials	5.4 mL
37.5 mg/m <sup>2</sup> (50 %)	67.5 mg	1 vial	2.7 mL
25 mg/m <sup>2</sup> (33 %)	45 mg	1 vial	1.8 mL

Manufactured by Baxter Oncology GmbH, Halle/Westfalen, Germany

Marketing Authorization Holder:  
Celgene Europe B.V., Netherlands

**Last revised in 07/2018**

#### To Report Any Side Effect(s):

##### *Saudi Arabia:*

– The National Pharmacovigilance and Drug Safety Centre (NPC)

- Fax: +966-11-205-7662
- Call NPC at +966-11-2038222, Exts: 2317-2356-2340
- Toll free phone: 8002490000
- E-mail: npc.drug@sfd.gov.sa
- Website: www.sfd.gov.sa/npc

##### *United Arab Emirates:*

Pharmacovigilance & Medical Device section  
P.O.Box: 1853  
Tel: 80011111  
Email: pv@moh.gov.ae  
Drug Department  
Ministry of Health & Prevention  
Dubai

##### *Oman:*

Department of Pharmacovigilance & Drug Information  
Directorate General of Pharmaceutical Affairs & Drug Control  
Ministry of Health, Sultanate of Oman  
Phone Nos. 22357687 / 22357686  
Fax: 22358489  
Email: dg-padc@moh.gov.om  
Website: www.moh.gov.om

##### • *Other Countries:*

– Please contact the relevant competent authority.

### This is a medicament

- A medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament to you
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not, by yourself, interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep Medicament out of the reach and sight of children and any other person not directly involved in the treatment

Council of Arab Health Ministers  
Union of Arab Pharmacists