

COQUELUSEDAL CHILDREN, suppository

Grindelia (soft extract) / Gelsemium (soft extract)

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

You must always take this medicine strictly according to the information provided in this leaflet or by your doctor or pharmacist.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, even not listed in this leaflet, tell your doctor or pharmacist (section 4).
- You must contact a doctor if your symptoms do not improve or worsen after 5 days.

1. WHAT COQUELUSEDAL CHILDREN IS AND WHAT IT IS USED FOR

COQUELUSEDAL CHILDREN is a traditional herbal medicine used in acute benign bronchial diseases in children from 30 months to 15 years old. Its use is reserved for specified indications exclusively based upon long-standing use.

2. BEFORE YOU USE COQUELUSEDAL CHILDREN

Do not use COQUELUSEDAL CHILDREN in the following cases:

- Allergy to the active substances, to other members of the Asteraceae (Compositae) family or any of the constituents of this medicine mentioned in section 6,

- Child under 30 months old,
- Child with a recent history of lesions of the anus or rectum, contraindication due to the route of administration.

Special warnings

In the event of emergence of dyspnea (breathlessness), fever or purulent sputum, ask your doctor for advice.

If the symptoms persist or worsen during the treatment, contact a doctor.

In case of chronic inflammation of the bronchi and the lungs, ask your doctor before using COQUELUSEDAL CHILDREN.

It contains 15 mg of propylene glycol per dose.

Intake or use of other medicines

Tell your doctor or pharmacist if you are taking or have recently taken or might use any other medicines.

Pregnancy breast-feeding and fertility

If you are pregnant, breast-feeding, if you think you are pregnant or if you plan a pregnancy, ask your doctor or pharmacist for advice before taking this medicine. In the absence of sufficient data, the use during pregnancy and lactation is not recommended.

3. HOW TO USE COQUELUSEDAL CHILDREN

Always take this medicine strictly according to the information provided in this leaflet or by your doctor or pharmacist. Ask your doctor or pharmacist in case of doubt.

Dosage:

- child from 30 months to 15 years old: 1 or 2 suppositories daily.
- child under 30 months old: this medicine is not indicated.

Administration route: rectal route.

Treatment duration: 5 days. If the symptoms persist after 5 days of treatment, consult a doctor or pharmacist.

If you forget to take COQUELUSEDAL CHILDREN, do not take a double dose to make up for a forgotten dose.

If you have other questions about the use of this medicine, ask more information to your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, COQUELUSEDAL CHILDREN can cause side effects, although not everybody gets them.

Local effects (irritation of the anus or rectum), related to the administration route, may occur. They are more frequent and severe, the longer the duration of use, the higher the frequency of administration and the higher the dosage.

Reporting side effects

If you feel any side effects, talk to your doctor or pharmacist. This also applies to any possible side effects not listed in this leaflet.

You can also report side effects directly through the national reporting system. By reporting side effects, you are helping to provide more information about drug safety.

5. HOW TO STORE COQUELUSEDAL CHILDREN

Keep out of the reach and sight of children. Store at a temperature not exceeding 30°C. Do not use after the expiry date stated on the box. The expiry date refers to the last day of the month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

Active substances:

Grindelia (*Grindelia robusta* Nutt.) (soft extract of flower top): 20 mg, extraction solvent: ethanol 70% V/V, drug-to-extract ratio 4/1.

Gelsemium (*Gelsemium sempervirens* (L) Ait.) (soft extract of underground part): 10 mg, extraction solvent: ethanol 60% V/V, drug-to-extract ratio 10/1, per 1 suppository.

Other ingredients: propylene glycol, hard fat.

This medicine is in box of 10 suppositories of brown green color, smooth in blister PVC/PE.

Chaoul Pharmaceuticals (CHA-PHA) S.A.L.
Lebanon (Packing)
Under License of ELERTE
Manufactured by Laboratoires ELERTE - France