

## Package leaflet: Information for the patient

# Carlevo

## 10/100 mg, 25/100 mg & 25/250 mg Tablets

### Carbidopa/Levodopa

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Carlevo is and what it is used for
2. What you need to know before you take Carlevo
3. How to take Carlevo
4. Possible side effects
5. How to store Carlevo
6. Contents of the pack and other information

#### 1. What Carlevo is and what it is used for

Carlevo improves the signs of Parkinson's disease. Parkinson's disease is a long-term illness where:

- you become slow and unsteady
- your muscles feel stiff
- you may develop shaking or trembling (called 'tremor').

If not treated, Parkinson's disease can make it hard for you to continue your normal daily activities.

Carlevo contains two different medicines called: levodopa and carbidopa.

- levodopa turns into a material called 'dopamine' in your brain. The dopamine helps to improve the signs of your Parkinson's disease.
- carbidopa belongs to a group of medicines called 'aromatic amino acid decarboxylase inhibitors'. It helps levodopa work more effectively by slowing the speed at which levodopa is broken down in your body.

#### 2. What you need to know before you take Carlevo

##### Do not take Carlevo:

- If you are allergic to **carbidopa or levodopa** or any of the other ingredients of this medicine (listed in section 6)
- If you have ever had skin cancer or you have any unusual moles which have not been examined by your doctor
- If you are taking certain medicines called 'MAOIs' (Monoamine Oxidase Inhibitors) used for depression. You need to stop using these medicines at least two weeks before you start Carlevo (see also under 'Other medicines and Carlevo' below)
- If you have a condition called 'narrow-angle glaucoma' that may cause a sudden build up of pressure in the eye
- If you have a severe mental disorder
- If you are pregnant, might become pregnant, or are breast-feeding.

Do not take Carlevo if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Carlevo.

#### Warnings and precautions

Talk to your doctor or pharmacist before taking Carlevo:

- If you have a history of fits (convulsions)
- If you have had an ulcer in your gut (called 'duodenal' or 'peptic ulcer') or have vomited blood
- If you have had a heart attack, heart beat problems, circulation or breathing problems
- If you have had kidney, liver or hormonal problems
- If you have had depression or other mental problems
- If you have a condition called 'chronic wide-angle glaucoma' that may cause a build up of pressure in the eye. You will need to have regular checks on the pressure in your eye.
- If you sometimes have sudden sleep attacks or sometimes feel very sleepy
- If you are due to have surgery.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Carlevo.

Tell your doctor if you or your family/carer notices you are developing addiction-like symptoms leading to craving for large doses of Carlevo and other medicines used to treat Parkinson's disease.

Tell your doctor if you or your family/carer notices you are developing urges or cravings to behave in ways that are unusual for you or you cannot resist the impulse, drive or temptation to carry out certain activities that could harm yourself or others.

These behaviours are called impulse control disorders and can include addictive gambling, excessive eating or spending, an abnormally high sex drive or an increase in sexual thoughts or feelings. Your doctor may need to review your treatments.

#### Other medicines and Carlevo

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. This includes herbal medicines. This is because Carlevo can affect the way some other medicines work. Also some other medicines can affect the way Carlevo works.

In particular tell your doctor or pharmacist if you are taking any of the following medicines:

- Medicines for Parkinson's disease containing levodopa:
  - If they are 'slow release', you will need to wait 24 hours after your last dose before starting Carlevo.
  - If they are 'normal release', you will need to wait 12 hours after your last dose before starting Carlevo.
- Tell the doctor or pharmacist even if you have only taken them in the past.
- Medicines for Parkinson's disease which do not contain levodopa will usually be continued. However, your dose may be changed.
- Medicines for mental problems (including depression), tuberculosis (TB), high blood pressure, muscle spasms, epilepsy or other diseases related to involuntary movements. Your dose may need to be changed.
- Medicines to treat low iron. Your dose may need to be changed.
- Medicines called monoamine oxidase inhibitors 'MAOIs' (see also '**Do not take Carlevo**').
- Anticholinergic medicines (such as orphenadrine, trihexyphenidyl, benztropine and procyclidine). Your dose may need to be changed.
- Phenytoin which is used to treat fits (convulsions).
- Papaverine which is used to treat impotence in men.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking this medicine. Your doctor or pharmacist has a more complete list of medicines to avoid while taking Carlevo.

#### Tests while you are taking Carlevo

This medication can affect some laboratory tests that your doctor may perform on blood or urine samples. Please remind your doctor if you are taking Carlevo and are having any tests.

#### Carlevo with food and drink

Try to avoid taking your tablets with a heavy meal. If your diet contains too much protein (meat, eggs, milk, cheese) Carlevo may not work as well as it should.

#### Pregnancy and breast-feeding

Do not take Carlevo if you are pregnant, might become pregnant or are breastfeeding. Levodopa, one of the substances in Carlevo, is passed into human milk.

Ask your doctor or pharmacist for advice before taking any medicine, if you are pregnant or breast-feeding.

#### Driving and using machines

- Carlevo affects different people in different ways. Some people have side effects which affect their ability to drive or use tools or machines (see Section 4 Possible side effects). Do not drive or use tools or machines if you get these effects.
- Carlevo can also make you sleepy or cause 'sudden sleep attacks'. If this happens to you, you must not drive or use tools or machines. Your doctor will tell you if you can start driving again if these attacks stop.

#### 3. How to take Carlevo

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

##### Taking this medicine

- Take this medicine by mouth.
- Although your medicine can have an effect after one day, it may take up to seven days to work.
- Take them at regular time intervals according to your doctor's instructions.
- Do not change the times at which you take your tablets or take any other medicines for Parkinson's disease without first consulting your doctor.
- Try to avoid taking your tablets with a heavy meal.

##### If you have not had levodopa before

The usual starting dose:

- for Carlevo 25 mg/100 mg tablets: one tablet three times a day.
- for Carlevo 10 mg/100 mg tablets: one tablet, three or four times a day.

##### If you have had levodopa before

Your doctor will ask you to stop taking your medicine for Parkinson's disease before you start taking Carlevo.

The usual starting dose is:

- for Carlevo 25 mg/100 mg tablets and Carlevo 25 mg/250 mg tablets: one tablet three or four times a day.
- for Carlevo 10 mg/100 mg tablets it will depend on what you were taking before.

More than one Carlevo product may be prescribed by your doctor. If you have been given different Carlevo tablets to take make sure that you are taking the correct one at the right time.

#### Children and adolescents (under 18 years of age)

Carlevo is not suitable for children under the age of 18 years.

#### If you take more Carlevo than you should

If you take too many tablets see your doctor immediately.

#### If you forget to take Carlevo

Do not take a double dose to make up for a forgotten dose.

#### If you stop taking Carlevo

Do not stop taking Carlevo or change your dose without talking to your doctor first. When you stop taking Carlevo the following can occur: stiff muscles, high temperature (fever) and mental changes.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

##### Stop taking Carlevo and see your doctor straight away, if you notice any of the following side effects:

- allergic reaction, the signs may include hives (nettle rash), itching, rash, swelling of your face, lips, tongue or throat. This may cause difficulty in breathing or swallowing
- chest pain
- uneven (irregular) heart beat or palpitations
- dizziness on standing-up quickly
- bleeding from your gut which may be seen as blood in your faeces or darkened faeces (gastro-intestinal bleeding)
- blood problems, the signs may include pale skin (pallor), tiredness, fever, sore throat or mild bruising and prolonged bleeding after injury
- stiff muscles, high fever
- mental changes including delusions, hallucinations and depression
- fits (convulsions).

##### The most common side effects are

- abnormal movements such as twitching or spasms (which may or may not be like your Parkinson's symptoms)
- nausea

##### Other side effects include

- fainting, anorexia, high blood pressure
- inflammation of the veins, being sick (vomiting), diarrhoea, discoloration of urine, sweat or saliva
- on-off phenomenon, characteristic of some people with long-standing Parkinson's disease. This is when you can have unpredictable changes from being mobile - "on" - to a sudden inability to move - "off". "Off" to "on" can occur just as suddenly.
- dizziness; sleepiness (including excessive drowsiness or sudden sleep onset episodes), pins and needles
- dream abnormalities, confusion, feeling agitated, shortness of breath, hair loss

##### You may experience the following side effects:

- Craving for large doses of Carlevo in excess of that required to control motor symptoms, known as dopamine dysregulation syndrome. Some patients experience severe abnormal involuntary movements (dyskinesias), mood swings or other side effects after taking large doses of Carlevo.
- Inability to resist the impulse to perform an action that could be harmful, which may include:
  - strong impulse to gamble excessively despite serious personal or family consequences.
  - altered or increased sexual interest and behaviour of significant concern to you or to others, for example, an increased sexual drive.
- uncontrollable excessive shopping or spending.
- binge eating (eating large amounts of food in a short time period) or compulsive eating (eating more food than normal and more than is needed to satisfy your hunger).

**Tell your doctor if you experience any of these behaviours; he/she will discuss ways of managing or reducing the symptoms.**

##### Side effects that have been reported with medicines containing levodopa.

These may be experienced when taking Carlevo.

##### Nervous system:

- loss of control over the voluntary movements of everyday life
- numbness, increased hand tremor, muscle twitching, muscle cramp, irregular movement of jaw muscles resulting in difficulty opening the mouth
- difficulty sleeping, feeling anxious or high, falling over and abnormal walking patterns
- headache

##### Eyes:

- drooping eyelid and dilated pupil
- changes in vision, irregular movement of the eye

##### Digestive system:

- indigestion, dry mouth, bitter taste
- swelling of the salivary glands, difficulty swallowing, grinding of the teeth
- hiccups, abdominal pain and distress, constipation, wind
- burning sensation of the tongue.

##### Sexual:

- persistent abnormal erection of the penis

##### Urinary:

- difficulty passing urine or incontinence (inability to control urine flow)

##### Skin:

- changed patches of pigmented skin, including, irritated or irregular moles, or moles in which you have noticed changes (melanoma)

##### General:

- weight gain or loss, swelling in the limbs
- flushing, hot flushes, increased sweating
- feeling weak, faint or tired
- hoarseness, general feeling of being unwell
- increased energy or activity, unusual breathing pattern

If any symptoms persist or you experience any other side effects please tell your doctor or pharmacist. It will help if you make a note of what you experienced, when it started and how long it lasted.

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. By reporting side effects you can help provide more information on the safety of this medicine.

#### 5. How to store Carlevo

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the blister and the outer packaging. The expiry date refers to the last day of that month.
- Do not store above 30°C. Keep away from humidity.
- Do not use this medicine if you notice visible signs of deterioration.
- Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

#### 6. Contents of the pack and other information

##### What Carlevo contains

- The active substances in Carlevo 25/100 mg are 25 mg of anhydrous carbidopa and 100 mg of levodopa.
- The active substances in Carlevo 10/100 mg are 10 mg of anhydrous carbidopa and 100 mg of levodopa.
- The active substances in Carlevo 25/250 mg are 25 mg of anhydrous carbidopa and 250 mg of levodopa.
- The other ingredients in Carlevo tablets are microcrystalline cellulose, pregelatinized starch, and magnesium stearate. Carlevo 10/100 mg and 25/250 mg tablets also contain FD&C Blue No. 2 (Aluminium Lake). Carlevo 25/100 mg tablets contain Spectracol brown oxide.

##### What Carlevo looks like and contents of the pack

- Carlevo 10/100 mg tablets are presented as blue, round tablets scored on one side.
- Carlevo 25/100 mg tablets are presented as brown, oval tablets scored on one side.
- Carlevo 25/250 mg tablets are presented as blue, oval tablets scored on one side.
- Carlevo 10/100 mg, Carlevo 25/100 mg and Carlevo 25/250 mg tablets are supplied in blister packs of 30, 100 and 20 tablets respectively, packed in carton box with a leaflet.

Not all strengths may be marketed.

##### Marketing Authorisation Holder and Manufacturer

Pharmaline s.a.l. – Lebanon  
P.O. Box: 90201 Jdeidet-El-Metrn, Lebanon  
Contact us: pharmaline@maliagroup.com  
Website: www.pharmaline.com.lb

Carlevo 10 mg/100 mg tablets Reg.N° Lebanon: 144720/1

Carlevo 25 mg/100 mg tablets Reg.N° Lebanon: 144720/1

Carlevo 25 mg/250 mg tablets Reg.N° Lebanon: 144820/1

##### This is a medicament:

- A medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.
- Keep all medicaments out of reach of children.

Council of Arab Health Ministers  
Union of Arab Pharmacists

**This leaflet was last revised in October 2021**

Rev. 10/2021 905376-A