

Fucidin® cream

Composition

Each gramme contains:
Fusidic acid 20 mg in neutral cream base.

Properties

Fucidin® is an antibiotic with powerful antibacterial activity against a number of gram-positive organisms. Staphylococci, including the strains resistant to penicillin or to other antibiotics, are particularly susceptible to Fucidin®.
Besides pronounced antibacterial activity against the majority of organisms responsible for skin infections, Fucidin® also possesses unique ability for penetration to the focus of infection, even when applied to intact skin.

Indications

Fucidin® cream is indicated for treatment of skin infections caused by staphylococci, streptococci, Propionibacterium acnes, Corynebacterium minutissimum, and other organisms sensitive to fusidic acid. The most important indications being:

| | |
|-----------------|----------------|
| Impetigo | Boils |
| Carbuncles | Paronychia |
| Infected wounds | Sycosis barbae |
| Hidradenitis | Erythrasma |
| Folliculitis | Acne vulgaris |

Fucidin® Cream is an invisible, non-staining preparation, which is cosmetically acceptable for treatment of face and scalp infections.

Application

Fucidin® Cream is applied to the affected area 2-3 times daily, generally for a period of 7 days, except of acne where therapy should be extended for a longer period according to the nature of this disease.

Side-effects

Fucidin® Cream is remarkably well tolerated, and there is an extremely low frequency of hypersensitivity reactions.

Storage conditions

Store at controlled room temperature (15°C-25°C).

Date of last revision of the package insert leaflet:

September 2008

LEO Pharmaceutical Products
Ballerup - Denmark

THIS IS A MEDICAMENT

- Medicament is a product, but not like others products.
- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the Doctor's prescription, the method of use and the instructions of the Pharmacist who sold you the medicament. The Doctor and Pharmacist are experts in medicaments, their benefits and their risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same medicament and do not increase doses without consulting your Doctor.

Do not leave medicament within reach of children

Council of Arab Health Ministers
Union of Arab Pharmacists



LEO

024236-00