

# MICOGYL

## VAGINAL SUPPOSITORIES

### COMPOSITION

Each vaginal suppository contains:  
Metronidazole 500 mg, Miconazole  
nitrate 100 mg, Excipients: Hard Fat q.s.

### PHARMACOLOGICAL PROPERTY

Combined preparation for intravaginal uses.  
It possesses antibacterial, antiprotozoal and  
antifungal activity.

Metronidazole is an antibacterial and  
antiprotozoal agent, active against anaerobic  
bacteria (*Bacteroides* spp., *Fusobacterium*  
spp., *Clostridium* spp., *Gardnerella vaginalis*,  
*Helicobacter pylori*, *Eubacteria*, and anaerobic  
cocci), and also protozoal infections  
(*Trichomonas* spp., *Giardia lamblia*,  
*Entamoeba histolytica*). It possesses a  
bactericidal mechanism of action.

Miconazole has a fungicidal action. It is  
particularly active against *Candida* spp.,  
*Trichophyton* spp., *Epidermophyton* spp.,  
*Microsporum* spp., *Pityrosporum* spp.

### ADMINISTRATION AND DOSAGE

Insert one vaginal suppository in the morning  
and at night for 7 days or insert one vaginal  
suppository at night for 14 days.

### ACTION AND USES

Micogyl is used in the treatment of candidal  
vulvovaginitis due to *Candida albicans*, in  
bacterial vaginosis due to anaerobic bacteria  
and *Gardnerella vaginalis*, in trichomonal

vaginitis due to *Trichomonas vaginalis* and in  
mixed vaginal infections.

### SIDE EFFECTS

Patients may experience local mild burning,  
itching or irritation immediately after inserting  
the vaginal suppository, abdominal cramps and  
headaches. Rarely hypersensitivity reactions  
have occurred.

### CONTRAINDICATIONS

Known hypersensitivity to Metronidazole  
and Miconazole nitrate or to any of the  
components of the vaginal suppositories.

### SPECIAL PRECAUTIONS:

It is recommended to avoid sexual contacts  
during the treatment. The base contained  
in the suppository formulation may interact  
with certain rubber or latex products, such  
as those used in vaginal contraceptive  
diaphragms or condoms, therefore,  
concurrent use is not recommended.

### USE IN PREGNANCY AND LACTATION

Use during pregnancy and lactation should  
be carefully considered. Micogyl should be  
administered only if evidently necessary.

### STORAGE

Store at temperature not exceeding 25°C.  
Keep out of the reach of children.