

RozyCal® D3

COMPOSITION

Calcium carbonate.....1500mg
(Equivalent to Calcium 600mg)

Cholecalciferol.....10mcg
(Equivalent to 400 IU of Vitamin D3)

Excipients: Corn Starch, Isomalt, Croscarmellose Sodium, Microcrystalline Cellulose, Magnesium Stearate, Sucralose, Povidone K30, Orange Flavor.

PHARMACEUTICAL FORM & PRESENTATION

Tablets pack of 60.

ROZYCAL®D3 IS INDICATED IN THE FOLLOWING CONDITIONS

- Prevention of calcium deficiency in adults & elderly.
- In cases of calcium and Vitamin D deficiency in adults and elderly.
- As an adjuvant to osteoporosis treatment if needed.

ROZYCAL®D3 IS NOT RECOMMENDED IN THE FOLLOWING CONDITIONS

- Hypersensitivity to Vitamin D or to any component.

PRECAUTIONS

- In case of simultaneous treatment with sodium fluoride or biphosphonate, it is advised to keep an interval of 2 hrs between doses, and in case of treatment with an antibiotic of the family of tetracyclines, it is advised to keep an interval of 3 hrs between doses.
- Consult your physician if you have renal insufficiency prior to taking any OTC product.

INTERACTIONS OF ROZYCAL®D3 WITH OTHER DRUGS

Consult your physician if you are being treated with digitalics, cyclines, biphosphonates, estramustine and iron salts while taking Rozy-cal®D3.

PREGNANCY AND LACTATION

Pregnant & Lactating women should consult their physician before taking any over the counter (OTC) product.

MODE OF ADMINISTRATION

1-2 tablets per day to be taken with a glass of water.

UNDESIRABLE EFFECTS

Calcium intake may cause in some individuals constipation or diarrhea, bloating, or nausea.

STORAGE INSTRUCTIONS

Store below 30°C protected from humidity. Keep in original pack in intact conditions.

Date of revision: August 2015.



Benta S.A.L.

Dbayeh- Lebanon