

**Denk
Pharma**

DEXA-TRESORTIL

Capsules

Please read carefully

Each capsule contains:

Indomethacin 25 mg

Dexamethasone 0,25 mg

Indications

Inflammatory rheumatic diseases such as rheumatoid arthritis, ankylosing spondylitis and gout. Inflammatory phases in osteoarthritis, disk lesions, degenerative joint disease. Inflammatory conditions such as bursitis, tendinitis and tenosynovitis.

Contra-indications

Active peptic ulcer or a history of recurrent gastro-intestinal; ulcerative colitis. Haemophilia and other bleeding disorders. Allergy to nonsteroidal antiinflammatory drugs (salicylates, etc).

The use of indomethacin is not recommended during pregnancy or lactation. It should not be administered to children.

Adverse effects

Headache, dizziness, light-headedness, confusion and other reactions from the central nervous system may occur and generally disappear with continued treatment or upon reduction of the dosage. Dyspepsia, gastrointestinal bleeding and ulceration, fluid retention, disturbed renal or hepatic function, fatigue, tinnitus, drowsiness or insomnia may occur.

Anaphylactic reactions in hypersensitive patients.

Interactions with other medications

Serum levels of indomethacin are raised by the concurrent administration of probenecid. Indomethacin reduces the natriuretic effect of frusemide.

Dosage and administration

Unless otherwise prescribed by the physician 2-6 capsules are administered daily in divided doses with meals. Maintenance dosage: 3 capsules. The capsules should be taken with some food, milk or antacid to avoid gastrointestinal disturbances.

The duration of treatment should be determined by the physician.

Storage

Store in a cool and dry place.

Do not use this product after the expiry date.

Keep out of reach of children.

Sole agents:

SADCO S.A.R.L.

E. DENK OHG, 81675 MUENCHEN
Prinzregentenstr. 79

Made in Germany

Vers. 003237/1