

Disflatyl®

Against flatulence and meteorism

Digestive disorders tend to be linked with the excessive accumulation of gas in the stomach and intestine. The most common symptoms developing through this are a sensation of pressure, tension and fullness in the epigastrium, eructations, flatus, pain and even intestinal spasm.

Being greatly distended, the intestinal loops may also force the diaphragm upwards mechanically, causing cardiac symptoms and dyspnoea (gastrocardiac syndrome). Symptoms like this do considerably interfere with general well-being and they are very common. Disflatyl is indicated with them and has proved effective.

Disflatyl offers a new way of treating such abnormal accumulation of gas in the gastrointestinal tract. Disflatyl contains activated dimethylpolysiloxane, a substance with surface action which causes the gas bubbles covered with mucus in the stomach and intestine to collapse.

Disflatyl brings rapid and noticeable relief of symptoms and removes the gas causing the flatulence. The most important active constituent of Disflatyl, dimethylpolysiloxane, is not absorbed from the stomach and intestine, so that it has no harmful effect whatever on the organism.

Composition

1 tablet contains:

Silicium dioxide	2 mg
Activated dimethylpolysiloxane	40 mg

Indications

Accumulation of gas in the gastrointestinal tract, sensation of pressure and fullness in the epigastrium, temporary meteorism following dietary indiscretions or lack of physical exercise, postoperative meteorism; gastrocardiac syndrome; in preparation for X-ray examination (stomach, intestine, gall bladder, kidney) and before gastroscopy.

Dosage

1 – 2 tablets to be thoroughly chewed after meals; in preparation for X-rays, give for 2 or 3 days before examination.

Packing

Box of 30 tablets

Box of 100 tablets

