

Concor® COR 2.5 mg



Active substance: Bisoprolol fumarate (2:1)

Composition

One film-coated tablet contains
Medicinally active ingredient:
2.5 mg bisoprolol fumarate (2:1).
Other ingredients:
Colloidal silicon dioxide, magnesium (stearate, palmitate, oleate), crospovidone, corn starch, microcrystalline cellulose, calcium hydrogen phosphate, dimethicone, macrogol 400, titanium dioxide (E171), hypromellose.

Presentation and package sizes

Film-coated tablet, off-white, heart-shaped, with dividing score
Packs of 30 tablets

Substance group

Cardioselective beta-receptor blocker

Marketing authorization holder and manufacturer

Merck KGaA, Frankfurter Strasse 250,
64293 Darmstadt, Germany

Indications

Treatment of stable chronic, moderate to severe heart failure (myocardial insufficiency) with impaired systolic ventricular function (ejection fraction ≤ 35 %, determined by echocardiography) in addition to ACE inhibitors and diuretics, and optionally cardiac glycosides.

Contraindications

When must you not take Concor COR 2.5 mg?

- Concor COR 2.5 mg must not be taken in:
- acute myocardial insufficiency (heart failure) or during deterioration (decompensation) of heart failure requiring intravenous therapy with substances increasing the contractility of the heart.
 - shock induced by disorders of cardiac function (cardiogenic shock)
 - severe disturbances of atrioventricular impulse conduction (second or third degree AV block) without a pacemaker
 - sick sinus syndrome
 - disturbed impulse conduction between sinus node and atrium (sinoatrial block)
 - markedly slowed heart beat (pulse rate less than 60 beats/min) before the start of treatment
 - markedly decreased blood pressure (systolic blood pressure below 100 mm Hg)
 - severe bronchial asthma or severe chronic obstructive lung disease
 - late stages of peripheral arterial occlusive disease or vascular spasms in toes and fingers (Raynaud's syndrome)
 - untreated tumours of the adrenal medulla (phaeochromocytoma)
 - metabolic acidosis
 - known hypersensitivity to bisoprolol or to any of the tablet constituents.

When may you take Concor COR 2.5 mg only after consulting your doctor?

In the following cases you may take Concor COR 2.5 mg only under certain conditions and with particular caution: Please ask your doctor about this. This also applies if you have been affected by any of the following in the past

- Bronchospasm (bronchial asthma, obstructive airways diseases)
- treatment with inhalation anaesthetics
- diabetes mellitus with extremely fluctuating blood glucose levels; symptoms of markedly reduced blood glucose (hypoglycaemia) can be masked
- strict fasting
- ongoing desensitisation therapy
- mild disturbances of atrioventricular impulse conduction (first degree AV block)
- disturbed cardiac blood flow due to vasospasms of the coronary vessels (Prinzmetal's angina)
- peripheral arterial occlusive disease (intensification of complaints may occur especially when starting therapy)

In patients with a personal or family history of psoriasis, beta-receptor blockers (e.g. Concor COR 2.5 mg) should only be used if the benefit-to-risk ratio has been carefully weighed.

In patients with a tumour of the adrenal medulla (phaeochromocytoma) Concor COR 2.5 mg may only be administered after previous alpha-receptor blockade.

What must pregnant women and nursing mothers pay attention to?

During pregnancy Concor COR 2.5 mg should only be used after the doctor has carefully weighed the benefit-to-risk ratio. In general beta-blockers reduce placental blood flow and can affect the development of the unborn child. Placental and uterine blood flow as well as the growth of the unborn child must be monitored and, if required, alternative therapeutic measures considered. The newborn child must be monitored closely after delivery. Symptoms of reduced blood glucose and slowed pulse rate generally occur within the first 3 days of life. It is not known whether bisoprolol passes into breast milk. Therefore, breastfeeding is not recommended during Concor COR 2.5 mg therapy.

What must be especially observed in children and elderly people?

Concor COR 2.5 mg should not be used in children as no adequate therapeutic experience in this context is available. No dose adjustment is required in elderly patients. No adequate therapeutic experience is available to date with regard to heart failure patients over 80 years of age.

Precautions for use and warnings

What precautions must be observed?

The treatment of myocardial insufficiency with Concor COR 2.5 mg requires regular monitoring. This is absolutely necessary especially at the start of therapy.

In bronchial asthma or other chronic obstructive pulmonary diseases that may be associated with symptoms concomitant bronchodilator therapy is indicated. An increase in airways resistance may occasionally occur in asthma patients, requiring a higher β2-sympathomimetic dose.

Like other beta-blockers, bisoprolol can increase the sensitivity to allergens and the severity of anaphylactic reactions, i.e. acute general allergic reactions. Adrenaline does not always produce the desired therapeutic effect in these cases.

The symptoms of thyroid hyperfunction (thyrotoxicosis) can be masked by bisoprolol.

So far no adequate therapeutic experience is available for Concor COR 2.5 mg in heart failure patients with concomitant insulin-dependent type-I diabetes mellitus, impaired kidney function (serum creatinine ≥ 3.4 mg/dl), impaired liver function, restrictive cardiomyopathy, congenital heart diseases or haemodynamically relevant cardiac valve diseases.

No adequate therapeutic experience is available either in patients with mild heart failure (NYHA II) as well as heart failure and myocardial infarction within the last 3 months. Therapy with bisoprolol should not be discontinued abruptly without compelling indication.

What must you pay attention to if you are driving a vehicle, operating machinery, or working without a firm hold?

In a study on patients with coronary heart disease bisoprolol did not affect the driving performance of the patients. However, due to individually different reactions to the drug, the ability to drive a vehicle, to operate machinery, or to work without a firm hold may be impaired. This is particularly the case at the start of treatment, when the dosage is increased or the medication changed, as well as in conjunction with alcohol.

Special warnings

THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of the treatment prescribed.
- Do not repeat the same prescription without consulting your doctor.

Keep medicament out of reach of children.

Council of Arab Health Ministers
Union of Arab Pharmacists

Drug interactions

What other drugs influence the effect of Concor COR 2.5 mg or what influence does Concor COR 2.5 mg have on the effect of other drugs?

Simultaneous administration of the following drugs is not recommended:

Increased lowering of blood pressure, delayed atrioventricular impulse conduction as well as reduced contractility of the heart muscle have been observed after simultaneous use of calcium antagonists.

Concurrent use of bisoprolol and clonidine can lead to a stronger reduction of heart rate and to delayed impulse conduction in the heart. Discontinuation of clonidine can also bring about an excessive increase in blood pressure.

Simultaneous administration of monoamine oxidase inhibitors (except MAO-B inhibitors) can affect the blood pressure (lowering of blood pressure, but also excessive increase in blood pressure).

The following drugs may only be used at the same time under certain conditions and with particular caution:

The depressant effects of Concor COR 2.5 mg and antiarrhythmics on impulse conduction and contractility of the heart can become additive. Parasympathomimetics (including tacrine) can prolong the atrioventricular conduction time.

Other beta-blockers, even if contained in eye drops, potentiate the effect of Concor COR 2.5 mg.

Concomitant use of Concor COR 2.5 mg and insulin or other drugs lowering the blood glucose level (oral antidiabetics) can potentiate the effect of the latter. Warning signs of reduced blood glucose (hypoglycaemia) – especially accelerated pulse (tachycardia) – can be masked or suppressed.

The concurrent administration of Concor COR 2.5 mg and anaesthetics may lead to a pronounced drop in blood pressure. Counter-regulatory mechanisms, e.g. increase in heart rate (reflex tachycardia) can be impaired. Continuation of beta blockade reduces the risk of arrhythmia during initiation of anaesthesia and intubation. The anaesthetist should be informed about treatment with Concor COR 2.5 mg. Concurrent therapy with Concor COR 2.5 mg and digitalis can lead to marked slowing of the heart beat and of impulse conduction in the heart. Simultaneous administration of mefloquine also furthers a slowing of the

heart beat. Prostaglandin synthesis inhibitors (e.g. acetylsalicylic acid) can reduce the antihypertensive effect of Concor COR 2.5 mg.

Concurrent administration of ergotamine derivatives (e.g. ergotamine containing drugs against migraine) and Concor COR 2.5 mg can lead to an increase in peripheral circulatory disturbances. Concomitant administration of Concor COR 2.5 mg and sympathomimetics can reduce the effect of the two substances. The treatment of allergic reactions may require an increased adrenaline dose.

The antihypertensive effect of Concor COR 2.5 mg can be potentiated by tricyclic antidepressants, barbiturates, phenothiazines and other antihypertensive drugs.

Concomitant administration of rifampicin and Concor COR 2.5 mg can slightly shorten the half-life of bisoprolol. Dose adjustment is in general not required.

Dosage instructions, mode and duration of administration

The following dosage instructions apply for Concor COR 2.5 mg, unless your doctor has prescribed otherwise. Please follow these directions carefully, otherwise Concor COR 2.5 mg cannot have the proper effect.

How many Concor COR 2.5 mg tablets should you take and how often should you take them?

You should have stable chronic heart failure without any acute deterioration (decompensation) within the last 6 weeks.

You should already be treated with an ACE inhibitor at optimal dosage (or, in the case of intolerance of ACE inhibitors, a different vasodilator), a diuretic and, if required, a digitalis preparation. This first-line medication should have remained largely unchanged for 2 weeks before starting therapy with Concor COR.

Treatment with bisoprolol should be initiated with slow, gradual dose increases. The attending physician should have experience in the therapy of chronic heart failure. Concor COR 2.5 mg is intended for initial treatment.

However, in some patients it may be sufficient for maintenance therapy. The following dose increases are recommended:

1.25 mg bisoprolol (equivalent to ½ tablet Concor COR 2.5 mg) once daily for 1 week.

If this dose is well tolerated increase to 2.5 mg bisoprolol (equivalent to 1 tablet Concor COR 2.5 mg) once daily for 1 week.

If this dose is well tolerated increase to 3.75 mg bisoprolol (equivalent to 1½ tablet Concor COR 2.5 mg) once daily for 1 week.

If this dose is well tolerated increase to 5 mg bisoprolol once daily for 4 weeks.

If this dose is well tolerated increase to 7.5 mg bisoprolol once daily for 4 weeks.

If this dose is well tolerated increase to 10 mg bisoprolol once daily as a maintenance dose.

For dosages over 2.5 mg bisoprolol correspondingly higher-strength tablets of Concor COR are available.

At the start of treatment with 1.25 mg bisoprolol (equivalent to ½ tablet Concor COR 2.5 mg) the patients should be monitored for 4 hours (blood pressure, heart rate, impulse conduction disorders as well as symptoms of heart failure).

The recommended maximal dose of 10 mg bisoprolol per day is reached at the earliest after a dose increase over 12 weeks and should not be exceeded.

A result of adverse reactions (e.g. slowing of heart beat with symptoms or drop in blood pressure or symptoms of deteriorating heart failure) may be that not all patients can be treated with the highest recommended dosage. If necessary, the beta-blocker dosage can be gradually reduced again, or the treatment (see below) can be discontinued and resumed again at a later time.

If deterioration of heart failure or intolerance occur during the titration phase the attending physician is advised to reduce or, in the presence of compelling reasons, immediately discontinue bisoprolol.

Treatment of stable chronic heart failure generally means long-term therapy.

Dosage in impaired liver and/or kidney function

Dose titration in heart failure patients with impaired liver or kidney function should be carried out with particular care since no pharmacokinetic investigations are available on this subject.

How and when should you take Concor COR 2.5 mg?

You should take the tablet whole with some liquid in the morning before, during or after breakfast.

How long should you take Concor COR 2.5 mg?

Treatment of stable chronic myocardial insufficiency is a long-term therapy. The dosage may only be changed by direction of your doctor. Also, treatment with Concor COR should only be interrupted or discontinued prematurely by direction of your doctor. For termination of therapy with Concor COR the dose should be gradually reduced (e.g. halving of the dose at weekly intervals).

You should not discontinue treatment with Concor COR abruptly as this may lead to temporary deterioration of the heart failure.

Incorrect intake and overdosage

What must you do if you have taken too many Concor COR 2.5 mg tablets?

In the case of suspected Concor COR 2.5 mg overdosage please inform your doctor immediately. Depending on the extent of overdosage your doctor can then decide which measures to take. The most frequent signs of Concor COR 2.5 mg overdosage include slowed heart beat (bradycardia), marked drop in blood pressure, bronchospasm, acute heart failure as well as hypoglycaemia.

In the case of overdosage therapy with Concor COR 2.5 mg should be discontinued after consultation of the attending physician.

What must you pay attention to if you have taken too little Concor COR 2.5 mg or have forgotten to take it?

Do not take the double dose the next time but continue your regimen as prescribed either under "Dosage instructions" or by your doctor.

What must you pay attention to if you interrupt treatment or stop it prematurely?

Please do not interrupt or stop treatment with Concor COR 2.5 mg without having consulted your doctor.

You should not discontinue treatment with Concor COR 2.5 mg abruptly as this may lead to temporary deterioration of the heart failure. The dose should be gradually reduced (e.g. halving of the dose at weekly intervals). This should be observed especially in patients with disturbed coronary blood flow.

Adverse effects

Besides the desired main effects, drugs can also have undesirable effects, so-called adverse reactions, which by no means occur in every patient, however.

What adverse reactions may occur when you are taking Concor COR 2.5 mg?

The following data are based on the therapeutic experience obtained after approval of bisoprolol for the therapy of hypertension as well as coronary heart disease.

- Central nervous system:

- | | |
|---------------|---|
| Occasionally: | Tiredness*, exhaustion*, dizziness*, headache* |
| Rare: | Sleep disorders, depression |
| Very rarely: | Nightmares, hallucinations |
| - Eyes: | |
| Very rarely: | Reduced tear flow (to be taken into consideration in patients wearing contact lenses) |
| Single cases: | Conjunctivitis |
| - Ears: | |
| Very rarely: | Hearing disorders |

- Cardiovascular system:

- | | |
|---------------|--|
| Occasionally: | Sensation of cold and numb extremities |
| Rare: | Slowing of heart beat (bradycardia), disturbed atrioventricular impulse conduction (AV block), deterioration of heart failure, stronger drop in blood pressure (also when standing up from a supine position, orthostatic hypotension) |

- Respiratory system:

- | | |
|--------------|--|
| Rare: | Bronchospasm in patients with a history of bronchial asthma or obstructive airways disease |
| Very rarely: | Allergic cold (rhinitis) |

- Gastrointestinal tract:

- | | |
|---------------|---|
| Occasionally: | Nausea, vomiting, diarrhoea, constipation |
|---------------|---|

- Muscles and skeleton (locomotor system):

- | | |
|-------|--------------------------------|
| Rare: | Muscle weakness, muscle cramps |
|-------|--------------------------------|

- Skin:

- | | |
|---------------|--|
| Very rarely: | Hypersensitivity reactions (itching, temporary flush, rash) |
| Single cases: | Beta-receptor blockers can trigger psoriasis, aggravate the condition or lead to psoriasiform rash. Loss of hair |

- Genito-urinary organs:

- | | |
|--------------|-------------------|
| Very rarely: | Potency disorders |
|--------------|-------------------|

- Metabolism:

- | | |
|--------------|--|
| Very rarely: | Increased liver enzyme levels (GOT, GPT), hepatitis, increased triglyceride levels |
|--------------|--|

*These symptoms occur especially at the start of treatment. They are mild and usually disappear within 1 to 2 weeks after the start of treatment. *If you should experience any adverse reactions not referred to in this package leaflet, you should report these also to your doctor or pharmacist.*

What countermeasures are to be taken in the case of adverse reactions?

Your doctor will decide about any possibly required countermeasures.

Notes on stability

The expiry date of this pack is printed on the folding box and on the brim of each blister strip. Do not use this pack once the expiry date has elapsed.

How should Concor COR 2.5 mg be stored?

Do not store Concor COR 2.5 mg above 25°C.

Date of issue

February 2002

Reg. № Lebanon 23649

Packed by Pharmaline - Lebanon
Licensed by Merck KGaA, Darmstadt, Germany