

Thilo-Tears Gel

Active constituents -2-

1 g gel contains 3 mg carbomer, 50 mg mannitol, 40 µg thiomersal for preservation, and 945.712 mg water for injection.

Indications -4-

"Dry eye" conditions, tear substitute for all conditions with deficient or defective wetting of the ocular surface.

Contraindications 5

Hypersensitivity against thiomersal.

Side effects 6

Occasionally temporary visual disturbances until the gel is evenly distributed over the ocular surface.

Dosage and administration 7

Instil one drop 4 times daily or more frequently into the conjunctival sac or as required to relieve the symptoms. The tube must be held vertically, so that the drop is released by gravity. This technique ensures that the size of the drop does not exceed the capacity of the conjunctival sac and that the eye-lashes do not stick together.

Thilo-Tears Gel is suitable for long-term use. Contact-lens users should not use Thilo-Tears Gel while wearing their lenses.

Special warnings 8

Thilo-Tears Gel should not be used after the expiry date printed on the label.

Thilo-Tears Gel should not be used for more than 6 weeks after the tube has been opened.

Store at room temperature.

Medicines should be kept out of the reach of children.

For more information, please consult the doctor or pharmacist.

Dr. Thilo & Co.
GmbH
D-7800 Freiburg



Thilo

Thilo-Tears Gel

35275-A

Administration of Thilo-Tears Gel

Instructions for Use

Thilo-Tears Gel

Dosage and administration

Instill one drop into the eye more than once a day. The conjunctival sac of the eye required to relieve the symptoms. The tube must be held vertically so that the drop is released by gravity. This technique ensures that the size of the drop does not exceed the capacity of the conjunctival sac and that the eye-lashes do not stick together.

Directions for use

A correct therapeutic drop-size is obtained if the tube is held

Properties

Thilo-Tears Gel forms a lubricating and protecting film on the cornea. It is employed as a tear substitute. In the treatment of severe cases of "dry eye" conditions, Thilo-Tears Gel has been shown to be especially useful, since the gel moisturizes the cornea and conjunctiva for a prolonged time, so that it has to be applied less frequently than conventional tear substitutes.

Active constituents: 1 g gel contains 10 mg thiomersal, 50 mg mannitol, 9.45 mg water for injection. Indications: "Dry eye" conditions, tear substitute for all conditions with deficient or defective wetting of the ocular surface.

vertically over the eye.

How supplied

10 g Thilo-Tears Gel

Dr. Thilo & Co.
GmbH
D-7800 Freiburg



Thilo