



# Travogen®

Bayer Schering Pharma

## This is a medicament

- ▶ A medicament is a product which affects your health and its consumption contrary to instructions is dangerous for you.
- ▶ Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- ▶ The doctor and the pharmacist are experts in medicine, its benefits and risks.
- ▶ Do not by yourself interrupt the period of treatment prescribed.
- ▶ Do not repeat the same prescription without consulting your doctor.

Keep medicament out  
of reach of children

Council of Arab Health Ministers  
Union of Arab Pharmacists

## ▶ Composition

Isoconazole nitrate 1%.

Excipients: polysorbate 60, sorbitan stearate, cetyl stearic alcohol, white vaseline, liquid paraffin, purified water.

## ▶ Pharmaceutical form and contents

Cream, tube containing 20 g.

## ▶ Pharmacotherapeutic group

Topical antifungal.

## ▶ Manufacturer

Intendis Manufacturing S.p.A., Italy  
a division of

Bayer Schering Pharma AG, Germany

## ▶ Therapeutic indications

Superficial mycoses (dermatophytes, candidiasis, pityriasis versicolor), erythrasma.

## ▶ Contraindications

Confirmed or presumed hypersensitivity to the ingredients of the medicinal product or to other imidazole antifungal derivatives.

## ▶ Precautions for use

The product must not be used in pregnant women unless absolutely necessary and under direct medical supervision.

## ▶ Interactions

None reported.

## ▶ Special warnings

Use of products for topical use, particularly

if prolonged, can give rise to sensitisation phenomena. In this case, use should be interrupted and suitable therapy should be instituted.

In the event of application to the face, avoid ocular contact with Travogen.

## ▶ Dose, method and duration of administration

Except when prescribed otherwise, the dosage of Travogen cream is 2 applications daily. To ensure the success of treatment, it is necessary to observe a few hygiene instructions carefully, including drying the interdigital spaces thoroughly in the case of mycosis localised to the feet. Local treatment of fungal infections usually lasts 2-3 weeks and sometimes 4 weeks in stubborn cases, especially involving the interdigital spaces. More prolonged treatments are also possible on specific medical prescription. It is often advisable, in the case of interdigital lesions, to insert a gauze dressing soaked in Travogen cream between the toes.

## ▶ Undesirable effects

Irritant skin phenomena, which occur quite rarely. The occurrence during the treatment of any undesirable effect not described in this information leaflet should be reported promptly to the treating doctor or pharmacist.

Caution: do not use the medicine after the expiry date stated on the packaging.

Store below 30°C.

▶ **Date of approval by the Ministry of Health**  
March 2005