

# PANDERM

## Cream, Ointment

### Composition

#### PANDERM Cream

Each gram of the cream contains:

<i>Active ingredients:</i>	Nystatin	100,000 I.U.
	Neomycin (as sulphate)	2.5mg
	Gramicidin	0.25mg
	Triamcinolone acetonide	1mg

*Excipients:* Cetostearyl alcohol, white soft paraffin, liquid paraffin, cetomacrogol, purified water, methyl and propyl paraben, sodium phosphate, and propylene glycol.

#### PANDERM Ointment

Each gram of the ointment contains:

<i>Active ingredients:</i>	Nystatin	100,000 I.U.
	Neomycin (as sulphate)	2.5mg
	Gramicidin	0.25mg
	Triamcinolone acetonide	1mg

*Excipients:* Liquid paraffin and white soft paraffin.

### Properties

The following four basic therapeutic properties of **PANDERM** make the product useful and sufficient to treat minor to moderate inflammatory skin lesions: anti-inflammatory, antipruritic, antibacterial, and antifungal. Triamcinolone acetonide is a corticosteroid which produces marked anti-inflammatory, antipruritic, and anti-allergic effects when applied topically; nystatin is an antifungal agent characterized by a specific anticandidal activity; and the two antibiotics, neomycin and gramicidin, provide together antibacterial properties against a wide range of Gram-positive and Gram-negative micro-organisms.

### Indications

**PANDERM** is indicated in the treatment of inflammatory skin lesions particularly when candidal or bacterial infections are suspected.

Examples are the following:

- |                                   |                       |
|-----------------------------------|-----------------------|
| - Atopic eczema                   | - Seborrheic eczema   |
| - Nummular eczema                 | - Stasis eczema       |
| - Contact eczema                  | - Otitis externa      |
| - Post traumatic infective eczema | - Neurodermatitis     |
| - Follicular eczema               | - Anogenital pruritus |
| - Infantile eczema                | - Psoriasis           |

### Application

Apply a thin film to the affected area 2-3 times daily. For moist or weeping surfaces, the water-miscible cream is preferable, but for dry lesions, the more emollient ointment is often more satisfactory.

### Contraindications

**PANDERM** should not be applied to viral lesions of the skin particularly herpes simplex, varicella, and vaccinia. Likewise, it should be avoided in fungal lesions not susceptible to nystatin.

**PANDERM** is not intended for ophthalmic use nor should it be applied to the external auditory canal in patients with perforated ear drum.

### Precautions

Topical corticosteroids should not be used extensively by pregnant women. The same precaution applies to infants.

Use of occlusive dressings is not recommended since it may increase the risk of sensitivity reactions.

Systemic absorption of topical preparations in general is higher when applied to skin lesions due to extensive burns or ulcerations.

### Side Effects

**PANDERM** is well tolerated. Sensitivity reactions, such as burning sensation, itching, and irritation may occasionally be experienced. Secondary infections may occur with the prolonged use of topical corticosteroids.

### Presentation

Collapsible tube containing 15gm cream or ointment.

\* Store at room temperature not exceeding 25°C.

#### THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicines, their benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.
- Keep all medicaments out of the reach of the children.

Council of Arab Health Ministers,  
Union of Arab Pharmacists.

Any information? Call Our Toll Free No. (971) 800-4994



Produced by: **juphar**  
Gulf Pharmaceutical Industries,  
Ras Al Khaimah, U. A. E.

