

# Saridon

Trade Mark

## Pain tablets

<b>Composition</b>	250 mg paracetamol, 150 mg propyphenazone and 50 mg caffeine.
<b>Properties and effects</b>	«Saridon» is a combination of analgesic and antipyretic substances with a small quantity of caffeine. Its components are carefully balanced to give potent and rapid relief of pain which lasts for several hours.
<b>Indications</b>	Pain, e.g. headache, toothache, menstrual pain, migraine, neuralgia, neuritis, postoperative and rheumatic pain; febrile illnesses (influenza, etc.).
<b>Dosage and administration</b>	<i>Single dose:</i> adults: 1–2 tablets; children of school age: $\frac{1}{2}$ –1 tablet. This single dose may be repeated three times in 24 hours. The tablets may be taken whole and washed down with a large volume of water or other suitable liquid. Analgesics should not be taken for prolonged periods of time or in high doses without consulting a doctor.
<b>Restrictions on use</b>	«Saridon» is contraindicated in the presence of severe disorders of kidney function, congenital deficiency of glucose-6-phosphate-dehydrogenase (hemolytic anemia), acute hepatic porphyria and evidence of allergic sensitivity to the active ingredients of «Saridon». «Saridon» should not be used in infants and children until they are of school age. The established medical principle of prescribing medicaments in early pregnancy only when absolutely indicated should be borne in mind.
<b>Undesirable effects</b>	«Saridon» is usually well tolerated. In some cases occasional allergic skin reactions have been observed (skin rash, urticaria).
<b>Interactions</b>	With «Saridon», as with other analgesics, patients should avoid taking alcohol while under treatment, since the individual response cannot be foreseen.
<b>Stability</b>	This medicine should not be used after the expiry date (EXP) shown on the pack.
<b>Packs</b>	Tablets <span style="float: right;">10, 250</span>

### This is a medicament

- A medicament is a product which effects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep medicaments out of reach of children

Council of Arab Health Ministers  
Union of Arab Pharmacists

Made in Switzerland by  
**F. Hoffmann-La Roche Ltd, Basel**