



## Caplets Extra Pain Reliever

### Composition

Each caplet contains:

**Active ingredients** : Paracetamol 500mg.  
Caffeine 65mg.

**Excipients** : Starch, povidone, potassium sorbate,  
glycerol, cross-linked starch, cellulose,  
stearic acid, and talc.

### Properties

**ADOL Extra** provides "extra" pain relieving property by virtue of its components which combat pain symptoms in different ways. Paracetamol has a well-known analgesic and antipyretic effects and caffeine has been reported to potentiate the analgesic effect of paracetamol and to expedite its onset of action. **ADOL Extra** relieves pain of almost any origin and reduces elevated body temperature.

### Indications

**ADOL Extra** is indicated for the effective and prompt relief of headache, migraine, sore throat, sinusitis pain, toothache, backache, rheumatic and muscular pain, neuralgia and menstrual discomfort.

**ADOL Extra** also relieves discomfort and helps to reduce fever in colds and influenza.

### Dosage

2 caplets with water 3 to 4 times daily, not exceeding 8 caplets in 24 hours.

Not recommended for children below 12 years of age.

### Precautions & Warnings

Do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed by the physician.

As with any drug, pregnant women or nursing mothers are better to seek medical advice before taking this product.

Do not consume excessive amounts of tea or coffee together with **ADOL Extra** as this may make the patient tense and irritable.

### Side Effects

**ADOL Extra** is generally safe when therapeutic doses are not exceeded. Allergic reactions, including skin rash, may occur.

### Overdosage

Massive overdosage of paracetamol (exceeding 10g for adults or 150mg/kg body weight for children) may cause hepatic toxicity. Early signs include nausea, vomiting, abdominal pain, and increased sweating. Clinical evidence may not be apparent until 48-72 hours.

Overdosage should be treated promptly by gastric lavage followed by I.V. N-acetylcysteine or oral methionine, which might be repeated as needed, in the light of laboratory results.

### Drug Interactions

This product should not be taken concomitantly with anticoagulants or antidiabetics unless otherwise directed by a physician.

### Presentation

**ADOL Extra caplets:** Packs of 24 caplets.

\* Store at room temperature not exceeding 25°C, in a dry place.

#### THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicines, their benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.
- Keep all medicaments out of the reach of children.

Council of Arab Health Ministers,  
Union of Arab Pharmacists.

Any information? Call Our Toll Free No. (971) 800-4994



Produced by: **Julphar**  
Gulf Pharmaceutical Industries,  
Ras Al Khaimah, U. A. E.

