

### Precautions:

- It should be applied only to intact skin, not to broken skin or open wounds, it should not come into contact with the eyes or mucous membranes.
- If applied on large areas over a prolonged period of time, systemic side effects may appear.
- It should not be taken by mouth.

### Contraindications:

It is contraindicated in patients hypersensitive to Diclofenac, Acetylsalicylic acid, NSAIDs, Propylene glycol and Isopropanol.

### Drug Interactions:

No drug interactions have been reported yet.

### Side Effects:

- Sometimes, Pruritus, erythema, rash and stinging may occur.
- Photosensitivity reactions have been reported in isolated cases.

### Presentations:

**Locagel® gel:** Each gram contains Diclofenac Diethylamine equivalent to Diclofenac Sodium 10 mg in tubes of 30 g and 50g

**This is a Medicament-Keep medicaments out of the reach of children**

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.



**Hayat Pharmaceutical Industries Co. Ltd.**  
P.O.Box 1564  
Amman 11118 Jordan.

## Locagel®

Diclofenac diethylamine

### Properties:

**Locagel®** is a nonsteroidal anti-inflammatory drug with analgesic and antirheumatic properties.

When rubbed into the skin it gives a soothing and cooling effect.

It penetrates into the inflamed area eliciting a clinical response characterized by a decrease in inflammatory swelling. **Locagel®** also reduces tenderness and pain on movement.

### Composition:

Diclofenac diethylamine

**Excipients:** Cetomacrogol 1000, Carbomer 934 P, Diethanolamine, Isopropyl alcohol, Propylene glycol, Lavender, Purified water, Mineral oil, Caprylic/Capric Triglycerides (Medium chain Triglycerides).

### Indications:

**Locagel®** is indicated for local treatment of:

- Localized rheumatic diseases e.g. osteoarthritis of the spine and peripheral joints.
- Back pain.
- Localized forms of soft tissue rheumatism e.g. tenosynovitis, shoulder-hand syndrome and bursitis.
- Inflammation of the ligaments, tendons, muscles, and joints due to trauma e.g. sprains, strains, bruises.

### Dosage & Administration:

**Locagel®** should be applied 3-4 times daily to the affected sites and gently rubbed in.