

# SPASMO - DIGESTIN

Enteric Coated Tablets

Digestant, Choleric,  
Spasmolytic and  
Antiflatulent

## Composition :

Each tablet contains :

|                           |        |
|---------------------------|--------|
| Papain                    | 100 mg |
| Sanzyme 3500              | 36 mg  |
| Sodium dehydrocholate     | 10 mg  |
| Dicyclomine hydrochloride | 5 mg   |
| Simethicone               | 30 mg  |

## Properties :

**Spasmo - digestin** is a well balanced combination with specific digestive, choleric, spasmolytic and antiflatulent properties. **Spasmo - digestin** provides the body with papain, the proteolytic enzyme, in addition to sanzyme 3500 which is a multienzyme complex containing mainly protease, amylase, lipase and cellulase. These enzymes act upon food contents of the GIT and catalyze their conversion into simpler and easily digestible and absorbable components. **Spasmo - digestin** therefore brings to completion hydrolysis of proteins into peptones and small peptides; starch into dextrins and simpler sugars; fats into fatty acids and glycerol, and cellulose into absorbable matter. **Spasmo - digestin** also exerts choleric effect through sodium dehydrocholate which improves the flow of bile necessary for emulsification and digestion of fats. The spasmolytic effect of **Spasmo - digestin** is due to dicyclomine hydrochloride which rapidly and effectively relieves spastic pain of the GI musculature and intestinal hypermotility. The antimuscarinic and direct antispasmodic actions of dicyclomine hydrochloride account for its high efficacy in irritable bowel syndrome (spastic colon). The antiflatulent effect of **Spasmo - digestin** is due to simethicone, the powerful defoaming agent. Simethicone reduces surface tension of gas bubbles in the GIT causing them to coalesce and be easily expelled. At the same time, simethicone prevents formation of mucus - surrounded gas pockets in the GIT. These properties collectively make **Spasmo - digestin** the drug of choice for rapid relief of indigestion, GI spasms and flatulence.

## Indications :

- Dyspepsia, post-prandial discomfort, epigastric fullness and flatulence associated with functional indigestion due to digestive enzyme deficiency or imbalance.
- Chronic enteritis, gastritis and pancreatitis.
- Following surgery on the stomach, small intestine or pancreas, and postgastrectomy syndrome.
- Irritable bowel syndrome (spastic colon) with such symptoms as anorexia, intestinal spasm, flatulence, constipation or diarrhea.
- Loss of appetite.
- Anorexia nervosa.

## Dosage :

1-2 tablets to be taken immediately before each meal.

## Packing :

A box containing 30 enteric coated tablets.

## Storage :

Keep in a cool dry place.

Keep out of the reach of children.

PRODUCED BY



ALEXANDRIA