

RECTACURE®

(Tribenoside and Lidocaine)

PROPERTIES AND PHARMACOKINETICS

Rectacure gives prompt relief from the unpleasant symptoms of haemorrhoids (pain, burning sensation, itching, and skin tension) which are chiefly due to secondary inflammation in the anal region.

The therapeutic activity of Tribenoside is attributable to the reduction of capillary permeability, and improvement of vascular tone, and to its local anti-inflammatory properties and antagonistic action on a number of endogenous substances, which play a role as mediators in the development of inflammation and pain. Lidocaine helps to provide rapid symptoms relief.

Rectacure absorption is designed to exert a local action, systemic absorption of the active substance, Tribenoside, should be minimal.

INDICATIONS

Rectacure is indicated for external and internal haemorrhoids.

DOSAGE

The cream and / or one suppository should be applied morning and evening, until the acute symptoms have disappeared, the dose can subsequently be reduced to the application of cream- or one suppository- once a day.

CONTRAINDICATIONS

Rectacure is contraindicated with known hypersensitivity to any of its components.

PRECAUTIONS

In case of significant bleeding from the anal region or other unusual symptoms, a medical diagnosis should be asked.

Pregnancy and Lactation: Like any other drug, Rectacure should be used with caution during pregnancy, especially in the first three months. It is not known whether the active substance passes into the breast milk, the benefits for the mother must be weighted against the risk to the child.

SIDE EFFECTS

Rectacure is well tolerated; in rare cases, a slight burning sensation, pain, or increased intestinal motility have been reported following use of the cream or suppositories.

OVERDOSAGE

No reports available on acute or chronic overdosage. Overdosage should not occur if the preparation is used in accordance to instructions.

PRESENTATION

Rectacure cream : Tribenoside 5 gm and Lidocaine HCl 2 gm / 100 gm cream

Rectacure suppository : Tribenoside 400 mg and Lidocaine HCl 40 mg / suppository

THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep medicament out of the reach of children



Manufactured by:
Jazeera Pharmaceutical Industries
P.O.Box 106229, Riyadh 11666, Saudi Arabia.