



REPADERM[®] Gel
(Aescin and Diethylamine salicylate)

PROPERTIES

Repaderm Gel is an anti-swelling, anti-inflammatory drug with analgesic properties, intended for topical administration. Aescin in Repaderm Gel inhibits the accumulation of fluid in the tissues and decreases tissue swelling rapidly. Aescin also effectively combats inflammatory processes at the affected area. Salicylate in Repaderm Gel enhances the anti-inflammatory action and effectively relieves pain from the affected area.

Repaderm Gel is a non-greasy preparation, and penetrates rapidly and completely into the affected tissues, even without rubbing, and cools the affected area.

INDICATIONS

Repaderm Gel is indicated in the following cases:

- All localized swellings resulting from injury or from inflammatory lesions such as contusions, crush injuries, sprains, bruises, hematoma and tenosynovitis.
- Painful conditions of the vertebral column such as neck pain, lumbago, and sciatica.

DOSAGE AND ADMINISTRATION

Unless otherwise directed, spread a thin layer of Repaderm Gel onto the skin of the affected area once daily or more frequently. It is not necessary to rub the gel into the skin and it could be done if desired.

Repaderm Gel could be used, even over a longer period, until the complaints relieve.

CONTRAINDICATIONS

Hypersensitivity to this medicament.

PRECAUTIONS

This medicament should not be applied to broken skin, mucous membranes or skin areas exposed to radiotherapy. If skin is present with eczema or ulceration, it is advisable to apply the gel on surrounding skin only.

SIDE EFFECTS

This medicament is generally well tolerated. Side effects are infrequent and mild in nature.

Allergic skin manifestations may occur in very isolated cases.

PRESENTATION

Repaderm Gel 100gm : Aescin 1 g and Diethylamine salicylate 5 g.

Repaderm Gel 40gm : Aescin 0.4 g and Diethylamine salicylate 2 g.

THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep medicament out of the reach of children



Manufactured by :
Jazeera Pharmaceutical Industries
P.O.Box 106229, Riyadh 11666, Saudi Arabia.

