

PROPERTIES

Solvex is a mucolytic agent that is often administered to facilitate expectoration by reducing sputum viscosity.

INDICATIONS

Solvex is indicated for the treatment of acute and chronic respiratory tract disorders associated with Viscid Mucus.

DOSAGE AND ADMINISTRATION

Adults & children above 10 years : 8 mg three times daily.

Children:

5 - 10 years : 4 mg (5 ml) 3 times daily. 2 - 5 years : 2 mg (2.5 ml) 3 times daily. Under 2 years : 1 mg (1.25 ml) 3 times daily.

CONTRAINDICATIONS

Solvex should not be used in patients hypersensitive to Bromhexine.

PRECAUTIONS

Teratogenicity has not been reported in animal studies but, as with other drugs, it is advisable to avoid use during the first trimester of pregnancy. Solvex syrup contain aspartame and therefore care should be taken in phenylketonuria.

Drug Interactions:

Administration of bromhexine together with antibiotics (amoxicillin, cefuroxime, erythromycin, doxycycline) leads to higher antibiotic concentration in the lung tissue. No clinically relevant unfavourable interactions with other medications have been reported.

SIDE EFFECTS

Rarely mild gastro-intestinal side effects may occur. Allergic reactions including skin rashes, bronchospasm, angiooedema, and anaphylaxis have also been reported very rarely.

OVERDOSAGE

Overdose with Bromhexine has not been reported. However, if occur, symptomatic treatment is recommended.

PRESENTATION

Svrup

Solvex : Bromhexine HCl BP 4 mg / 5 ml.

Tablets

Solvex 4: Bromhexine HCl BP 4 mg / tablet.

Solvex 8: Bromhexine HCl BP 8 mg / tablet.

THIS IS A MEDICAMENT fedicament is a product which affects your health, and its consumption contrary

instructions is dangerous for you.

Follow strictly the doctor's prescription, the method of use and the instructions of pharmsoist who sold the medicament.

The doctor and the pharmacist are experts in medicine, its benefits a Do not by yourself interrupt period of treatment prescribed for you. Do not repeat the same prescription without consulting your doctor.

Keep medicament out of the reach of children

