# catarstat®

eve drops

PR	ES	EN'	TATI	ON .
----	----	-----	------	------

Eye drops : 10 ml bottle.

#### COMPOSITION .

ITION ;	
Active principles :	
Pyridoxine hydrochloride	122 00000000
Pyridoxine hydrochloride L-glutamic acid	0.02 g
Magnesium-potassium L-aspartate Glycine	0.38 g
Glycine	0.10 g
Preservative ·	0.24 g
Benzalkonium chloride Excipient : o s	Service Service
Excipient : q.s.	0.01 g
HITIC INCIDATIONS	100 ml

# THERAPEUTIC INDICATIONS :

This medication is for use as eye drops. It is intended to be used in lens opacities (cataract).

## CONTRAINDICATIONS :

IF YOU ARE IN ANY DOUBT, YOU SHOULD SEEK THE ADVICE OF YOUR PHYSICIAN OR PHARMACIST.

#### PRECAUTIONS .

IF YOU ARE IN ANY DOUBT. DO NOT HESITATE TO SEEK THE ADVICE OF YOUR PHYSICIAN OR PHARMACIST. IN ORDER TO AVOID POSSIBLE INTERACTIONS BETWEEN

SEVERAL MEDICATIONS YOU SHOULD SYSTEMATICALLY POINT OUT ANY OTHER PENDING TREATMENT TO YOUR PHYSICIAN OR PHARMACIST.

ALWAYS KEEP OUT OF REACH OF CHILDREN.

OTHER POSSIBLE EFFECTS OF THE MEDICATION:
AS IN THE CASE OF ALL ACTIVE SUBSTANCES, THIS MEDI-CATION MAY HAVE TROUBLESOME EFFECTS OF VARYING DEGREE IN CERTAIN INDIVIDUALS DO NOT HESITATE TO SEEK THE ADVICE OF YOUR PHYSICIAN OR PHARMACIST.

#### DOSAGE .

Generally 2 drops in each eye, 2 to 3 times daily. YOU SHOULD AT ALL TIMES STRICTLY FOLLOW YOUR PHY-SICIAN'S PRESCRIPTION.

### ADMINISTRATION :

See diagrams.

# PRESCRIBING CONDITIONS :

THIS MEDICATION HAS BEEN PRESCRIBED FOR YOU PER-SONALLY IN A SPECIFIC SITUATION

- IT MAY NOT BE APPROPRIATE IN OTHER SITUATION.

DO NOT SUGGEST THAT ANYBODY ELSE SHOULD USE IT.

#### SHELF-LIFE :

DO NOT EXCEED THE EXPIRY DATE CLEARLY SHOWN ON THE PACKAGE

# SPECIAL CONDITIONS OF CONSERVATION :

This medication should not be kept for more than 15 days after opening of the bottle.

