

EXOMUC[®] 200 mg

acetylcysteine

COMPOSITION

Acetylcysteine (INN)200 mg

Excipients: aspartam, beta-carotene, sorbitol, orange flavour q.s.p. one sachet.

Granules for oral solution in a sachet. Sugar-free.

THERAPEUTIC INDICATIONS

This medicine is an expectorant, it encourages coughing to remove mucus from the bronchi. It is recommended in the treatment of bronchial obstruction that accompanies certain conditions of the respiratory tract.

CONTRAINDICATION

This medicine **MUST NOT BE USED** in case of phenylketonuria (because of the presence of aspartam).

PRECAUTIONS

- Inform your doctor if you have a gastric or duodenal ulcer.
- It is advisable not to take an antitussive medication (cough suppressant) or a medicine that dries up bronchial secretions during the period that you are being treated with this medicine.

DRUG INTERACTIONS

SYSTEMATICALLY REPORT ANY OTHER ONGOING TREATMENT TO YOUR DOCTOR OR PHARMACIST SO AS TO AVOID ANY INTERACTION BETWEEN SEVERAL MEDICINES.

PREGNANCY AND BREAST-FEEDING

Inform your doctor if you are pregnant or breast-feeding.

IN GENERAL, YOU SHOULD ALWAYS ASK YOUR DOCTOR OR PHARMACIST FOR ADVICE BEFORE TAKING ANY MEDICINE IF YOU ARE PREGNANT OR BREAST-FEEDING.

DOSAGE AND METHOD OF ADMINISTRATION

Oral use.

Adults and children over 7 years of age : one sachet three times daily.

Children aged between 2 and 7 years : one sachet twice daily.

Number of calories per sachet : 3.5 Kcal.

Dissolve the granules in half a glass of water and then swallow.

UNDESIRABLE EFFECTS

LIKE ALL MEDICINE, EXOMUC MAY, IN CERTAIN PEOPLE, GIVE RISE TO VARYING DEGREES OF UNPLEASANT SIDE EFFECTS:

Possibility of gastrointestinal disturbances : nausea, diarrhoea, stomach pain. In such cases, it is recommended that the dosage be reduced.

Juillet 2007

007358 / V 38068-07

Laboratoires BOUCHARA - RECORDATI

68, rue Marjolin
92300 Levallois-Perret (FRANCE)

FABRICANT/MANUFACTURER/المُصنِّع:

SOPHARTEX
28500 Vernouillet (FRANCE)