

# CUSI ERYTHROMYCIN<sup>®</sup>

0,50% Eye ointment  
tube of 3.5g

**COMPOSITION:** Each g of ointment contains 5 mg of erythromycin.

Excipients: s.q.

**INDICATIONS:** Treatment of superficial ocular infections caused by microorganisms sensitive to erythromycin.

Prophylaxis of ophthalmia in the newborn.

**DOSAGE:** In the treatment of superficial ocular infections, normally apply once a day or more frequently, depending on the severity of the infection.

In the prophylaxis of ophthalmia in the newborn, a single application should be carried out.

## **CORRECT ADMINISTRATION PROCEDURE:**

**Application to the inside of the eye:** Separate the eyelids from the eye and apply a small amount (approximately a grain of rice) of ointment into the conjunctival sac.

**Application to the outside of the eye:** Soften and remove any scab with warm water, and apply the product directly onto the affected area.

Application of the ointment should be carried out under hygienic conditions.

Do not touch the tube tip to any surface.

Close tube after every application.

**CONTRAINDICATIONS:** Patients with hypersensitivity to erythromycin.

**ADVERSE REACTIONS:** Allergic reactions may occur which would require interruption of the treatment. The use of antibiotics may give rise to growth of non-susceptible microorganisms, particularly fungi, in this case medication should be interrupted and appropriate measures taken.

**INCOMPATIBILITIES:** Have not been described.

**OVERDOSAGE:** Due to the characteristics of this preparation, intended for topical use, no toxic effects are expected with the use of this product at the recommended. In the event of overdosage or accidental ingestion, consult physician or a Poison Control Center.

( **HOW SUPPLIED:** 3.5 g tube. )

~~KEEP ALL MEDICATION OUT OF THE REACH OF CHILDREN~~