

POLYDEXA®

Ear solution - Drops

COMPOSITION

Neomycin (INN) sulphate 1 g i.e. 650 000 U
Polymyxin B (INN) sulphate 1 000 000 U
Dexamethasone (INN) sodium metasulphobenzoate 0.100 g
Excipients: sodium mercuriothiolate, citric acid, sodium hydroxide, macrogol 400, polysorbate 80, water q.s. 100 ml. Bottle of 10.5 ml.

THERAPEUTIC INDICATIONS

This medicine is used in the treatment of intact eardrum external otitis, especially in case of infected eczema of the external auditory canal.

CONTRA-INDICATIONS

This medicine **MUST NOT BE USED** in case of :

- perforated eardrums,
- history of allergy to one of the ingredients.

WARNINGS

Do not swallow. Do not use without medical advice. As a precaution before any administration, the integrity of your eardrum should be verified by a physician. If the eardrum is perforated, this medicine may cause irreversible adverse effects in the ear (deafness, balance disorders).

PRECAUTIONS FOR USE

The duration of treatment must not exceed 10 days. If symptoms persist, consult your doctor.

INTERACTIONS WITH OTHER MEDICAMENTS AND OTHER FORMS OF INTERACTIONS

IN ORDER TO AVOID POSSIBLE INTERACTIONS WITH OTHER MEDICAMENTS, YOU MUST ALWAYS INFORM YOUR DOCTOR OR PHARMACIST OF ANY OTHER CURRENT TREATMENT.

PREGNANCY - LACTATION

This medicine can only be used during pregnancy or breastfeeding on the advice of your doctor. If you find out that you are pregnant during the treatment, consult your doctor who is the only person who can evaluate whether treatment can be continued.

POSODOLOGY AND METHOD OF ADMINISTRATION

Otic route. Comply with your doctor's prescription.

As an indication :

Adult : 1 to 5 drops, 2 times daily.

Child : 1 to 2 drops, 2 times daily.

In general, it is not recommended to use under pressure. Do not exceed 10 days of treatment. If symptoms persist, consult your doctor.

ADVERSE EFFECTS

Risk of local reactions (irritation) and antibiotic-related allergy.

In case of perforated eardrum, risk of deafness and balance disorders.

Possibility of selection of resistant bacteria and development of mycosis (infections due to fungi).

- Allergic reactions to the active ingredients.
- Local reactions (irritation) and antibiotic-related allergy.
- In case of perforated eardrum, risk of deafness and balance disorders.
- Possibility of selection of resistant bacteria and development of mycosis (infections due to fungi).

Mai 2002

V 58568-08

BOUCHARA-RECORDATI

68, rue Marjolin - 92300 Levallois-Perret (FRANCE)