

DIFRAREL® 100

Coated tablets

QUALITATIVE AND QUANTITATIVE COMPOSITION

ANTHOCYANOSIDIC EXTRACT OF VACCINIUM MYRTILLUS	100 mg
BETACAROTENE	5 mg

Excipient: Kieseluhr (diatomite for oenology); Starchy icing sugar, Talc, Corn Starch, Magnesium Stearate, Colophony, Turpentine type "Pinus pinaster", Shellac, Saccharose, Gelatine, Gum Arabic, Titanium Dioxide, Patent Blue V, Opaglos 600.

For one coated tablet.

PHARMACEUTICAL PRESENTATION

Box of 20 coated tablets.

PHARMACO-THERAPEUTIC CLASSIFICATION

VASCULAR PROTECTANT

(C: cardiovascular system)

(S: sensory organs)

NAME AND ADDRESS OF THE PRODUCER

Laboratoires LEURQUIN MEDIOANUM

68-88 rue Ampère - Z.I. des Chanoux

93330 NEUILLY SUR MARNE

Medical information : ☎ 01 49 44 29 29

CASE(S) IN WHICH THIS MEDICINE CAN BE USED

This medicine is recommended for the treatment of :

- venous circulation disorders,
- fragility of small blood vessels,
- impaired vision due to circulatory disorder.

WARNING!

CASE(S) IN WHICH THIS MEDICINE SHOULD NOT BE USED

This medicine SHOULD NOT BE USED in the following case:

- previous history of allergy to one of the constituents of the coated tablet;
- IF IN DOUBT, DO NOT HESITATE TO CONSULT YOUR DOCTOR OR CHEMIST.*

PRECAUTIONS

This medicine is fully effective when associated with a healthy lifestyle. Avoid exposure to sun, heat, standing for long periods, over-weight.

Walking and possibly wearing support stockings helps blood circulation.

If discomfort and fragility of blood vessels increase (spreading of bruising and ecchymosis), consult your doctor immediately.

IF IN DOUBT, DO NOT HESITATE TO CONSULT YOUR DOCTOR OR CHEMIST.

INTERACTION WITH OTHER MEDICINES AND OTHER INTERACTIONS

TO AVOID POSSIBLE INTERACTION BETWEEN SEVERAL MEDICINES, YOUR DOCTOR OR CHEMIST MUST BE SYSTEMATICALLY INFORMED OF ANY OTHER CURRENT TREATMENT.

PREGNANCY - BREAST FEEDING

It is not advisable to take this medicine during pregnancy.

If you should find that you are pregnant while taking the medicine, consult your doctor who can best decide on the necessity to continue.

Avoid taking this medicine when breast feeding.

Generally speaking, during pregnancy and while breast feeding, you should always consult your doctor or chemist before taking any medicine.

ADMINISTRATION OF THIS MEDICINE

DOSE

3 to 6 tablets a day.

The treatment is to be taken for 20 days per month, then renewed if necessary.

THIS MEDICINE HAS BEEN ISSUED OR PRESCRIBED FOR YOU PERSONALLY FOR A PARTICULAR REASON:

- *IT MAY NOT BE SUITABLE IN ANOTHER CASE,*
- *DO NOT ENCOURAGE USE BY ANOTHER PERSON.*

MODE AND METHOD OF ADMINISTRATION

To be taken orally.

Swallow the tablets, without chewing, with a glass of water.

LENGTH OF TREATMENT

The course of treatment should be taken for 20 days per month and renewed if necessary. Follow your doctor's recommendations.

UNDESIRABLE, TROUBLESOME SIDE-EFFECTS

AS IS THE CASE FOR ANY ACTIVE SUBSTANCES, THIS MEDICINE MAY, CAUSE UNDESIRABLE OR TROUBLESOME SIDE-EFFECTS IN CERTAIN INDIVIDUALS: digestive disturbances, in rare instances.

INFORM YOUR DOCTOR OR CHEMIST OF ANY UNDESIRABLE OR TROUBLESOME SIDE-EFFECT NOT MENTIONED ON THIS LEAFLET.

SPECIFIC PRECAUTIONS FOR STORAGE

To be stored at a temperature not exceeding 25°C.

STORAGE

DO NOT EXCEED THE EXPIRY DATE SHOWN ON THE OUTER PACKAGING.

LEAFLET REVISION DATE

August 24th 2005.