Anginovag aerosol

NEW ECOLOGIC CONTAINER: FOR THE PROTECTION OF THE ECOSYSTEM FREON GAS HAS BEEN ELIMINATED

DO NOT REMOVE THE CANNULA FROM THE BOTTLE ONCE ASSEMBLED AND USED

Pharmacological action

Anginovag combines in its formula active principles with antiinflamatory/analgesic and antibiotic actions. It is supplied in the form of spray with doping valve and that is why it is suitable for the preventive and curative therapy of the buccopharyngeal affections.

Composition

Each 100 ml contains: B-glycyrrhetinic acid, 60 mg; Degualinium chloride, 100 mg; Tyrothricin, 400 mg; Hydrocortisone acetate, 60 mg; Lidocaine hydrochloride, 100 mg; Excipients: (Sodium Saccharin) g.s. 100 ml.

Indications

Adjuvant treatment in prevention and healing of oropharyngeal disorders: Amygdalitis, pharyngitis, laryngitis, stomatitis, ulcers, aphthae, glossitis. Dosage

indicacion 8

Attack dose: 1-2 applications every 2-3 hours.

Maintenance or preventive dose: One application every 6 hours.

Contraindications/Side-effects/Interactions None are known.

Precautions

For its correct administration, it is recommended to place the flask in upright position, with the cannula pointing towards the place of application, but without introducing it completely into the mouth. Press the upper part of the cap from top to bottom up to the buffer.

Use during pregnancy It can be administered.

Overdosage and its treatment

Given the scarce toxicity of the preparation the appearance of sympthoms of poisoning is not foreseen, even in case of overdosage.

Cabron cacion y su tre into

Supply

Package containing 10 ml.

Keep out of sight and reach of children.

Manufactured by: FERRER INTERNACIONAL, S.A.