

# HEXALYSE®

## tablet to be sucked

### IDENTIFICATION OF THE MEDICINAL PRODUCT

#### **COMPOSITION**

BICLOTYMOL .....	5.0 mg
LYSOZYME HYDROCHLORIDE .....	5.0 mg
ENOXOLONE .....	5.0 mg

Excipients : Lemon essential oil, citric acid monohydrate, orange Yellow S (E 110), magnesium stearate, gum arabic, icing sugar for one tablet to be sucked.

#### **PHARMACEUTICAL FORM**

Tablet to be sucked, box of 30.

#### **PHARMACOTHERAPEUTIC GROUP**

FOR ANTI-INFLAMMATORY and ANTISEPTIC USE

(R: respiratory system).

#### **NAME AND ADDRESS OF THE MARKETING COMPANY**

BOUCHARA-RECORDATI Laboratories  
68, rue Marjolin - 92300 LEVALLOIS-PERRET - France

#### **WHEN SHOULD THIS MEDICINE BE USED?**

This medicine is recommended for treating mild sore throats in patients with no high temperature.

#### **WARNING!**

#### **WHEN NOT TO USE THIS MEDICINE?**

This medicine *SHOULD NOT BE USED* in the following circumstances:

- in children under 6 years of age,
- in patients with a history of allergy to biclotymol or to any of the other ingredients of the tablet.

#### **PRECAUTIONS FOR USE**

If the patient is diabetic or is on a low sugar diet, take quantity of sugar in each tablet into account in the daily allowance: 1.1 g.

If a high temperature and/or new problems should occur (significant sore throat or headache, purulent expectoration (sputum), difficulty in swallowing food...); **CONSULT YOUR DOCTOR.**

Do not use this medicine for an extended period of time.  
IF IN DOUBT, DO NOT HESITATE TO ASK YOUR DOCTOR OR  
PHARMACIST FOR ADVICE.

## **INTERACTIONS WITH OTHER MEDICINES AND OTHER INTERACTIONS**

This medicine contains bictotymol, lysozyme hydrochloride and enoxolone. Other medicines also contain these ingredients. Do not take them at the same time so as not to exceed maximum recommended doses (see dosage and method of administration).

*TO AVOID POSSIBLE INTERACTIONS BETWEEN DIFFERENT MEDICINES, YOU SHOULD ALWAYS TELL YOUR DOCTOR OR PHARMACIST ABOUT ANY OTHER TREATMENT YOU ARE TAKING.*

## **PREGNANCY - BREAST-FEEDING**

As a general rule, you should always seek the advice of your doctor or pharmacist before taking any medicine during pregnancy and lactation.

## **LIST OF EXCIPIENTS WHICH NEED TO BE KNOWN FOR SAFE USE IN CERTAIN PATIENTS**

Orange Yellow S (E 110).

## **HOW TO USE THIS MEDICINE**

### **DOSAGE**

Suck the tablets slowly, without chewing them.  
Six to eight tablets to be sucked every 24 hours, spreading out the doses.

*THIS MEDICINE HAS BEEN DISPENSED OR PRESCRIBED FOR YOU TO TREAT YOUR SYMPTOMS:*

- *IT SHOULD NOT BE GIVEN TO ANYONE ELSE.*
- *YOU SHOULD NOT RECOMMEND IT TO ANYONE ELSE.*

### **METHOD AND ROUTE OF ADMINISTRATION**

Suck the tablets slowly, without chewing them.

Oral route.

### **DURATION OF TREATMENT**

Do not use this medicine for an extended period of time.

### **UNDESIRABLE AND UNPLEASANT EFFECTS**

*AS WITH ALL ACTIVE PRODUCTS, THIS MEDICINE CAN LEAD TO ANNOYING EFFECTS OF VARYING SEVERITY IN SOME PEOPLE:*

*skin rash: if this occurs, stop taking the treatment.*

*INFORM YOUR DOCTOR OR PHARMACIST ABOUT ANY UNWANTED AND ANNOYING EFFECTS WHICH ARE NOT MENTIONED IN THIS LEAFLET.*

### **STORAGE**

**DO NOT USE AFTER THE EXPIRY DATE STATED ON THE OUTER PACKAGING**

### **SPECIAL STORAGE PRECAUTIONS**

Store at a temperature below 25°C.

### **DATE OF REVISION OF THE INFORMATION LEAFLET**

March 2005.