

# Hexaspray®

## Throat spray

Please read this entire leaflet carefully before you start taking this medicine.

- Keep this leaflet; you may need to read it again.

- If you have further questions, or any doubt, ask your doctor or pharmacist.

This medicine has been prescribed for you. You should not pass it on to others. It may harm them, even if their symptoms are the same as yours.

### IDENTIFICATION OF THE MEDICINAL PRODUCT

#### COMPOSITION:

Biclotymol ..... 750 mg  
Excipients: Benzyl alcohol, sodium edetate, methylparahydroxybenzoate (E218), anise oil, ammonium glycyrrhizinate, sodium saccharin, microcrystalline cellulose and carboxymethylcellulose, soybean lecithin, glycerol, ethylic alcohol, purified water s.q.f 30 g. Nitrogen (propulsive gas).

#### PHARMACEUTICAL FORM:

Throat spray. Pressurised bottle of 30 g.

#### PHARMACOTHERAPEUTIC GROUP:

Local antiseptic.

#### NAME AND ADDRESS OF THE MARKETING COMPANY:

BOUCHARA-RECORDATI Laboratories  
68 Rue Marjolin - 92 300 LEVALLOIS-PERRET - France

#### WHEN TO USE THIS MEDICINE?

This medicine is recommended for treating mild sore throats in patients with no high temperature.

#### WARNING!

##### WHEN NOT TO USE THIS MEDICINE?

This medicine MUST NEVER BE USED in the following situations:

- allergy to one of the ingredients of the throat spray
- throat sprays are contraindicated in children below 30 months of age (risk of laryngospasm).

**IF THERE IS ANY DOUBT, DO NOT HESITATE TO ASK YOUR DOCTOR OR PHARMACIST FOR ADVICE**

#### SPECIAL WARNINGS

KEEP OUT OF THE REACH OF CHILDREN.

Prolonged use (over 5 days) should ideally be avoided as it may alter the natural microbial balance in the throat.

The concomitant or successive use of antiseptics must be avoided given the potential interferences (antagonism, inactivation).

#### PRECAUTIONS FOR USE

If a high temperature, purulent excretions (sputum), difficulty in swallowing food occur, or if you feel worse or not better after 5 days: CONSULT YOUR DOCTOR.

#### INTERACTIONS WITH OTHER MEDICINES AND OTHER INTERACTIONS

*IN ORDER TO AVOID ANY POTENTIAL DRUG INTERACTION ALWAYS INFORM YOUR DOCTOR OR PHARMACIST OF ANY OTHER CONCOMITANT TREATMENT.*

This medicine contains biclotymol. Other medicines also contain this ingredient. Do not take them at the same time so as not to exceed the maximum recommended doses (see Dosage and method of administration).

#### PREGNANCY AND BREAST-FEEDING

##### Pregnancy

Ask your doctor or pharmacist for advice before taking this medicine.

Biclotymol should preferably not be administered during pregnancy. *IN GENERAL, IF YOU ARE PREGNANT OR BREAST-FEEDING, YOU SHOULD ALWAYS SEEK THE ADVICE OF YOUR DOCTOR OR PHARMACIST BEFORE TAKING ANY MEDICINE.*

**LIST OF EXCIPIENTS WHICH NEED TO BE KNOWN FOR SAFE USE IN CERTAIN PATIENTS:** Methyl parahydroxybenzoate (E218).

#### HOW TO USE THIS MEDICINE?

##### DOSAGE

**ONLY FOR ADULTS AND CHILDREN OVER 30 MONTHS.**

2 sprays 3 times per day.

#### METHOD AND ROUTE OF ADMINISTRATION

Oral route

##### SHAKE THE BOTTLE BEFORE EACH USE

Turn the spray system and press  
HOLD THE BOTTLE VERTICALLY



1  
Turn



2  
Press



YES



NO

#### DURATION OF TREATMENT

The treatment should not exceed 5 days.

#### UNWANTED AND UNPLEASANT EFFECTS

*LIKE ALL MEDICINES, THIS PRODUCT MAY PRODUCE UNPLEASANT EFFECTS OF VARYING SEVERITY IN SOME PEOPLE:*

Possibility of allergic symptoms: lip oedemas, urticaria, skin rashes, erythema and Quincke's oedema.

The presence of methyl parahydroxybenzoate (E218) and soya lecithin may trigger allergic reactions.

**TELL YOUR DOCTOR OR PHARMACIST ABOUT ANY UNWANTED OR UNPLEASANT EFFECT WHICH IS NOT MENTIONED.**

#### STORAGE

**DO NOT USE AFTER THE EXPIRY DATE SHOWN ON THE OUTER PACKAGING.**

#### SPECIAL STORAGE PRECAUTIONS

Keep away from heat.

#### DATE OF REVISION OF THE INFORMATION LEAFLET:

September 2005