

Bronquium

ELIXIR

A rational therapy for the treatment of respiratory obstruction

Pharmacological action. Bronquium Elixir contains theophylline having a bronchodilator action and glyceryl guaiacolate having an expectorant action. It is indicated for the treatment of respiratory affections where such actions are needed.

Composition

Each 100 ml contains: Anhydrous theophylline, 1 g;
Glyceryl guaiacolate, 0.6 g; Excipients, p.s.

Indications

Symptomatic treatment of bronchial asthma and other bronchospastic conditions such as asthmatic bronchitis and chronic bronchitis.

Dosage

Adults: 15-30 ml, 2-3 times daily.

Children: Weighing over 45 Kg: The same dose as the adults.

Over 8 years old: 15 ml, 2-3 times daily.

Under 8 years old: 2.5 ml per each 4.5 Kg body weight, 2-3 times daily.

During the first day of therapy and in severe attacks, the usual dose may be increased by one half.

Contraindications

Serious myocardic lesions.

Side-effects

Sporadically gastrointestinal disorders may appear or stimulating effects may occur (palpitations, insomnia, etc.).

Owing to individual idiosyncrasy undesirable actions may appear, and in such a case the treatment must be stopped.

Interactions

It must not be administered at the same times as preparations containing xanthine derivatives. The sympathicomimetic agents may increase the side-effects of theophylline.

Precautions / Warnings

The preparation must not be administered more frequently than every 6 hours.

After the administration by rectal route of any preparation containing theophylline or aminophylline, a period of 12 hours must elapse before administering the preparation.

Use during pregnancy

Administer with care and under medical vigilance.

Overdosage and its treatment

In case of poisoning by overdose proceed to gastric lavage and symptomatic treatment with sedatives, electrolytic sera and oxygen if necessary.

Supply

Bottle of 120 ml.

KEEP OUT OF REACH OF CHILDREN



FERRER INTERNACIONAL, S.A.
Gran Vía de Carlos III, 94
BARCELONA (España, Spain)