

intétrix®

1. IDENTIFICATION OF THE MEDICINAL PRODUCT

NAME

INTÉTRIX, capsule.

COMPOSITION:

TILQUINOL 50 mg
TILQUINOL laurylsulphate 50 mg
TILBROQUINOL 200 mg

Excipients : monohydrated lactose, corn starch.

For one capsule*

* Composition of the capsule cap: gelatine, titanium dioxide (E171), azorubine (E122), indigotine (E132).

PHARMACEUTICAL FORM:

Capsule, box of 20 or 40.

PHARMACO-THERAPEUTIC:

ANTI-PROTOZOAL

(P: Antiparasitic)

DISTRIBUTOR:

IPSEN Pharma
65, quai Georges Gorse 92100 Boulogne-Billancourt

MANUFACTURER:

BEAUFOUR IPSSEN Industrie rue Ethe Virton
28100 Dreux

2. WHAT IS YOUR MEDICINE USED FOR?

INTÉTRIX is recommended in the treatment of intestinal amoebiasis in adult.

3. WARNING !

WHEN YOU SHOULD NOT TAKE THIS MEDICINE:

IF YOU HAVE ANY DOUBT, YOU MUST CONSULT EITHER YOUR DOCTOR OR PHARMACIST.

PRECAUTIONS FOR USE:

Consult your doctor in case of hepatic insufficiency. In case of jaundice stop the treatment. INTÉTRIX must not be prescribed in association with other medicines containing hydroxyquinolines. Notify your doctor of any other ongoing treatment. *IF YOU HAVE ANY DOUBT, DO NOT HESITATE TO ASK YOUR DOCTOR OR PHARMACIST FOR ADVICE.*

INTERACTIONS WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION:

IN ORDER TO AVOID ANY POSSIBLE MEDICINAL PRODUCT INTERACTIONS, YOU SHOULD NOTIFY SYSTEMATICALLY YOUR DOCTOR OR PHARMACIST OF ANY OTHER ONGOING CONCOMITANT TREATMENT.

PREGNANCY AND BREAST-FEEDING

INTÉTRIX is not to be used during pregnancy.

If during the treatment you discover you are pregnant, you should quickly consult your doctor only he can adapt the treatment to you.

In general, during pregnancy and lactation, always ask your doctor or pharmacist for medical advice before taking any drug.

LIST OF EXCIPIENTS, WHICH MUST BE KNOWN FOR THE SAFE USE OF THE MEDICINAL PRODUCT IN SOME PATIENTS.

Azorubine (E122).

4. HOW TO USE THIS MEDICINAL PRODUCT?

DOSAGE:

2 capsules in the morning and 2 capsules in the evening.

METHOD AND ROUTE OF ADMINISTRATION:

Oral route.

FREQUENCY AND TIME AT WHICH THIS MEDICINAL PRODUCT SHOULD BE ADMINISTERED:

The product has to be taken at the beginning of meals.

DURATION OF TREATMENT:

The treatment should not exceed 10 days.

5. UNEXPECTED AND BOTHERSOME EFFECTS

LIKE ANY OTHER ACTIVE PRODUCT THIS MEDICINAL PRODUCT CAN, IN SOME PATIENTS, CAUSE A GREATER OR LESSER DEGREE OF DISCOMFORT.

– cutaneous reactions,

– exceptional nervous effects,

– biological effects: increase of transaminases (liver enzymes).

DO NOT HESITATE TO ASK YOUR DOCTOR OR PHARMACIST FOR ADVICE AND TO NOTIFY ANY UNEXPECTED AND BOTHERING EFFECT NOT MENTIONED IN THIS PACKAGE LEAFLET.

6. STORAGE

DO NOT EXCEED THE EXPIRY DATE PLAINLY INDICATED ON THE PACKAGING.

SPECIAL STORAGE PRECAUTIONS.

Should be stored at a temperature of less than 25°C.

7. REVISION DATE OF THE PACKAGE LEAFLET

June 1999.

