

Inongan

CREAM

NAME OF THE MEDICINAL PRODUCT: INONGAN, Cream

QUALITATIVE AND QUANTITATIVE COMPOSITION:

METHYL SALICYLATE: 12.50 g

RACEMIC CAMPHOR: 1.00 g

for 100 g of cream

EXCIPIENTS:

Sedefos 75® (ethylene distearate and diethylene glycol associated with polyoxyethylene fatty alcohol phosphate). Labrafil M2130® (saturated polyglycosylated glycerides), liquid paraffin, racemic menthol, eucalyptus oil, purified water.

PHARMACEUTICAL FORM: Cream, tube of 55 g.

PHARMACOTHERAPEUTIC CLASS:
analgesic (M: musculoskeletal system)

WHAT THIS MEDICINE IS USED FOR

Local symptomatic treatment of muscle, tendon and ligament pain in adults.

ATTENTION!

WHEN MUST THIS MEDICINE NOT BE USED

This medicine **MUST NOT BE USED** in the following cases:

- Allergy to salicylates or related drugs (non-steroidal anti-inflammatory drugs, aspirin) or to any of the ingredients.
- Do not apply on mucous membranes, in the eyes, on broken skin, on a moist or infected lesion, or under an occlusive bandage.

IF YOU ARE IN ANY DOUBT, YOU MUST CONSULT YOUR DOCTOR OR YOUR PHARMACIST FOR ADVICE.

SPECIAL WARNINGS:

This medicine contains camphor and menthol (terpene derivatives) which, at excessive doses, may cause convulsions in infants and children.

Follow the recommendations for use:

- Never apply more than the recommended dose (see Dosage section).
- Do not apply over a large area of the body.
- Follow the recommended frequency and duration of treatment.

PRECAUTIONS FOR USE:

- This medicine is restricted to adults.
- In case of previous or recent epilepsy, **ASK YOUR DOCTOR FOR ADVICE.**
- Wash your hands after use to avoid accidental contact with eyes and mucous membranes.

IF YOU ARE IN ANY DOUBT, DO NOT HESITATE TO SEEK THE ADVICE OF YOUR PHARMACIST OR YOUR DOCTOR.

Inongan

CREAM

INTERACTIONS WITH OTHER MEDICAMENTS AND OTHER FORMS OF INTERACTIONS

IN ORDER TO AVOID POSSIBLE INTERACTIONS BETWEEN VARIOUS DRUGS, YOU MUST ALWAYS TELL YOUR DOCTOR OR YOUR PHARMACIST ABOUT ANY OTHER TREATMENT YOU ARE RECEIVING.

PREGNANCY – BREASTFEEDING

If you are breastfeeding, it is preferable that you not use this medicine, and in no case should you apply it on the breasts.

As a general rule, if you are pregnant or breastfeeding you should always seek the advice of your doctor or pharmacist before taking a medication.

HOW TO USE THIS MEDICINE

DOSAGE: RESTRICTED TO ADULTS

Local massage on the painful area, one or more times a day.

Wash your hands after use.

METHOD AND ROUTE OF ADMINISTRATION: Cutaneous route.

DURATION OF TREATMENT: Do not use for prolonged periods without medical advice.

POSSIBLE SIDE EFFECTS:

Like all medicines, this medicinal product may cause more or less bothersome effects in some people:

- possible local allergic reactions requiring discontinuation of the treatment
- possible normal redness of the skin at the site of application
- if the recommended doses are not followed, possible agitation and confusion in the elderly (due to the presence of camphor and menthol).

DO NOT HESITATE TO ASK YOUR DOCTOR OR PHARMACIST FOR ADVICE AND TO TELL YOUR DOCTOR OR PHARMACIST IF YOU NOTICE ANY SIDE EFFECTS NOT LISTED IN THIS LEAFLET

STORAGE

DO NOT USE AFTER THE USE-BY DATE SHOWN ON THE OUTER PACKAGING

SPECIAL STORAGE PRECAUTIONS

Store at a temperature at or below 25°C.

MARKETING AUTHORISATION NUMBER: 341 589-4

MARKETING AUTHORISATION HOLDER:

SOFIBEL: 110-114 rue Victor Hugo – 92686 Levallois-Perret Cedex – France

DATE OF REVISION OF THE LEAFLET: September 1997