

DIOVENOR[®] 600mg

Diosmin film-coated tablet

Please read this leaflet carefully before taking this medicine. If you have any questions or doubts, ask for more information from your doctor or pharmacist. Keep this leaflet; you may need to refer to it again.

COMPOSITION: diosmin (quantity expressed in anhydrous and pure diosmin): 600 mg. Excipients: talc, anhydrous hydrophobic colloidal silica, micronized stearic acid, hypromellose, microcrystalline cellulose, macrogol 400 stearate, propyleneglycol, titanium dioxide, red cochineal A aluminium lacquer, black iron oxide, red iron oxide, carnauba wax, bee wax, gum lacquer, ethylic alcohol 95°. For one film-coated tablet. **PHARMACEUTICAL FORM:** film-coated tablets. **PHARMACO-THERAPEUTICAL GROUP:** VASCULOPROTECTOR/ACTING ON CAPILLARIES (C05CA03: cardiovascular tract). **IN WHICH SITUATIONS THIS DRUG MUST BE USED:** This drug is a venotonic (it increases the tonus of venous walls) and a vasculo-protector (it increases the resistance of the small blood vessels). It is preconised in: • blood circulation troubles (heavy legs, pain, restlessness when lying down). • Treatment of functional signs related to a haemorrhoidal attack. **IN WHICH SITUATIONS THIS DRUG MUST NOT BE USED:** known allergy to one of the components, this drug is generally contra-indicated during breastfeeding. **WARNINGS:** If troubles and fragility of vessels do not decrease within 15 days, consult your doctor. If haemorrhoidal symptoms persist after some days of treatment, it is advisable to consult your doctor. **PRECAUTIONS FOR USE:** This drug has the maximal efficiency when associated to healthy habits. Avoid sun, warm, prolonged standing, excess of weight. Walking and wearing appropriated stockings improve the blood circulation. **DRUG INTERACTIONS:** IN ORDER TO AVOID DRUG INTERACTIONS BETWEEN SEVERAL DRUGS INDICATE SYSTEMATICALLY ALL CURRENT TREATMENT TO YOUR DOCTOR OR YOUR PHARMACIST. **PREGNANCY:** This drug will be used during pregnancy only if necessary according to your physician's advice. If you discover that you are pregnant during the treatment ask to your doctor because he can judge if it is necessary to go on this treatment. **LACTATION:** In the absence of data concerning passage into breast milk, the treatment is not recommended during lactation. AS A GENERAL RULE IT IS ALWAYS ADVISABLE TO ASK YOUR PHYSICIAN OR YOUR PHARMACIST BEFORE USING A DRUG DURING PREGNANCY AND LACTATION. **LIST OF EXCIPIENTS WHICH HAVE TO BE WELL KNOWN TO ENSURE THE RISK-FREE USE OF THIS DRUG:** cochineal red A (E124). **HOW TO USE THIS MEDICINAL PRODUCT: DOSAGE:** • Venous insufficiency: 1 tablet per day, in the morning before breakfast. • Haemorrhoidal attack: 2 to 3 tablets per day, during meals. IN ALL CASES, RESPECT YOUR DOCTOR'S PRESCRIPTION. **WAY OF USE AND ROUTE OF ADMINISTRATION:** Oral route. Swallow the tablet with a glass of water. **DURATION OF TREATMENT, FREQUENCY AND MOMENT WHEN THE DRUG MUST BE TAKEN:** IN ALL CASES, RESPECT STRICTLY YOUR PHYSICIAN'S PRESCRIPTION. **SIDE EFFECTS:** AS FOR ALL DRUGS, SOME SIDE EFFECTS CAN OCCUR IN SOME PATIENTS: possibility of minor digestive troubles leading rarely to stop the treatment, due to presence of cochineal red A, risk of allergic reactions. INFORM YOUR DOCTOR OR YOUR PHARMACIST ABOUT ALL SIDE EFFECT NOT MENTIONED IN THIS LEAFLET. **STORAGE:** DO NOT EXCEED THE EXPIRY DATE CLEARLY PRINTED ON THE OUTER PACK. **DATE OF REVISION OF THE LEAFLET:** September 29, 2008. MA: 337 269-9

TITULAIRE/M.A. HOLDER :

LABORATOIRES INNOTHERA - 22, avenue Aristide Briand - 94110 Arcueil - France

FABRICANT/MANUFACTURER :

INNOTHERA CHOUZY - Rue René Chantereau - L'Isle Vert - 41150 Chouzy-sur-Cisse - France