

berialdistsc need for anti-inflammations vonant-siq ni eau to voldic

(Diclofenac diethylamine)

emulgel

Pharmacological properties

Voldic® emulgel (Diclofenac diethylamine) is a non-steroidal anti-inflammatory agent with a pronounced analgesic anti-rheumatic activity.

In cases of rheumatic inflammation or trauma, when **Voldic® emulgel** is applied topically, it relieves pain and gives soothing and cooling effects.

Indications

Usually **Voldic® emulgel** is indicated in the following :

- Localized soft tissue rheumatism: shoulder-hand syndrome, bursitis, tendovaginitis and periarthropathy.
- Localized rheumatic diseases: osteoarthritis and low back pain.
- Post-traumatic inflammation due to sprains, strains or bruises in the muscles, ligaments, and joints.

Dosage and administration

Usual adult dose:

Voldic® emulgel is applied 3-4 times daily to the affected area, and rubbed gently. The amount to be applied usually depends on the size of area to be treated, while the duration depends on the severity of the indication and the patient's response.

Voldic® emulgel can be combined with other forms of **Voldic®**

Usual children dose:

Use in children is not preferred, since no dose recommendation has been established.

Use in pregnancy and lactation

Safety for use during pregnancy and lactation has not been established.

Contraindications

Hypersensitivity to diclofenac or other NSAIDs, acetyl salicylic acid, and to propylene glycol or isopropanol.

Precautions

- Diclofenac diethylamine should be applied only to intact skin and not to open wounds.
- Contact with eyes or mucous membranes should be avoided.

Side effects

Generally, Voldic® emulgel is well tolerated, occasionally contact dermatitis of allergic or non allergic type with symptoms including itching, erythema, edema, rash, papules, vesicles, and scaling occurred. Photosensitivity reactions were also reported.

Systemic reactions are not likely to occur during topical treatment with Voldic® emulgel unless applied to large areas and for long periods of time.

Overdosage

In the case of accidental overdosage (e.g. in children), the usual measures employed to manage overdose with NSAIDs should be followed.

Presentation

Voldic® emulgel : Diclofenac diethylamine equivalent to 10 mg diclofenac sodium / g .(Tube containing 30 g) .

(This is a Medicament - Keep medicaments out of the reach of children)

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, method for use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.



Pharma International