

# AZONIT<sup>®</sup>

( Isoconazole nitrate )

## ***Pharmacological information***

**Azonit<sup>®</sup>** (isoconazole nitrate) is an antimycotic agent with a very broad spectrum antimicrobial activity. It is effective against yeasts, yeast-like fungi, dermatophytes and moulds. Also, **Azonit<sup>®</sup>** exerts activity against gram-positive bacteria.

## ***Indications***

- Treatment of superficial fungal infections.
- Pityriasis versicolor.
- Erythrasma.
- Vulvo – vaginal mycosis.
- **Azonit<sup>®</sup>** vaginal cream is also indicated in cases of fungal infections of the vagina.

## ***Dosage and administration***

**Azonit<sup>®</sup>** must be applied once daily. Hygienic measures should be employed for successful treatment. Also treatment should be continued for two to three weeks and up to four weeks particularly in the infections of the interdigital spaces.

To avoid recurrence, treatment should be continued for at least 2 weeks after clinical healing.

## ***Treatment of fungal infections of the vagina:***

**Azonit**® vaginal cream must be applied once daily using the applicator for seven consecutive days. The applicator must be inserted deep into the vagina. Treatment shouldn't be carried out during menstruation.

To avoid reinfection, concurrent treatment of the partner with **Azonit**® cream is recommended, all personal towels, underwear should be changed daily and boiled.

## ***Side effects***

**Azonit**® is well tolerated. The most encountered side effects, which are mild in nature are: skin or mucosal irritation and allergic skin reactions.

## ***Precautions***

Avoid contact with the eyes.

## ***Contraindications***

There are no known contraindications till now.

## ***Presentation***

**Azonit**® cream: 10mg Isoconazole nitrate/gm. (Tube contains 20 gm).

**Azonit**® vaginal cream: 10mg Isoconazole nitrate/gm. (Tube contains 40 gm).

( This is a medication - keep medications out of reach of children )



Pharma International

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, method for use and the instructions of the pharmacist who sold the medicament .
- The doctor and the pharmacist are experts in medicine, its benefits and risks .
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.