

Enclosure


[We note your agreement to the labeling below in an electronic communication dated November 15, 2002. Additionally, we note that, at this time, you intend to market only the 10 mg, 15 mg, 20 mg, and 30 mg dosage strengths. However, the Agency is approving all of the following dosage strengths: 2 mg, 5 mg, 10 mg, 15 mg, 20 mg, and 30 mg. Additionally, for completeness, we are including these dosage strengths into the labeling.]

ABILIFY[®] (aripiprazole) Tablets

Rx only

DESCRIPTION

ABILIFY[™] (aripiprazole) is a psychotropic drug that is available as tablets for oral administration. Aripiprazole is 7-[4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy]-3,4-dihydrocarbostyryl. The empirical formula is $C_{23}H_{27}Cl_2N_3O_2$ and its molecular weight is 448.38. The chemical structure is:



ABILIFY tablets are available in 2-mg, 5-mg, 10-mg, 15-mg, 20-mg, and 30-mg strengths. Inactive ingredients include lactose monohydrate, cornstarch, microcrystalline cellulose, hydroxypropyl cellulose, and magnesium stearate. Colorants include ferric oxide (yellow or red) and FD&C Blue No.2 Aluminum Lake.

CLINICAL PHARMACOLOGY

Pharmacodynamics

Aripiprazole exhibits high affinity for dopamine D_2 and D_3 , serotonin $5-HT_{1A}$ and $5-HT_{2A}$ receptors (K_i values of 0.34, 0.8, 1.7, and 3.4 nM, respectively), moderate affinity for dopamine D_4 , serotonin $5-HT_{2C}$ and $5-HT_7$, α_1 -adrenergic and histamine H_1 receptors (K_i values of 44, 15, 39, 57, and 61 nM, respectively), and moderate affinity for the

serotonin reuptake site ($K_i = 98 \text{ nM}$). Aripiprazole has no appreciable affinity for cholinergic muscarinic receptors ($IC_{50} > 1000 \text{ nM}$). Aripiprazole functions as a partial agonist at the dopamine D_2 and the serotonin $5-HT_{1A}$ receptors, and as an antagonist at serotonin $5-HT_{2A}$ receptor.

The mechanism of action of aripiprazole, as with other drugs having efficacy in schizophrenia, is unknown. However, it has been proposed that the efficacy of aripiprazole is mediated through a combination of partial agonist activity at D_2 and $5-HT_{1A}$ receptors and antagonist activity at $5-HT_{2A}$ receptors. Actions at receptors other than D_2 , $5-HT_{1A}$, and $5-HT_{2A}$ may explain some of the other clinical effects of aripiprazole, eg, the orthostatic hypotension observed with aripiprazole may be explained by its antagonist activity at adrenergic α_1 receptors.

Pharmacokinetics

ABILIFY activity is presumably primarily due to the parent drug, aripiprazole, and to a lesser extent, to its major metabolite dehydro-aripiprazole, which has been shown to have affinities for D_2 receptors similar to the parent drug and represents 40% of the parent drug exposure in plasma. The mean elimination half-lives are about 75 hours and 94 hours for aripiprazole and dehydro-aripiprazole, respectively. Steady-state concentrations are attained within 14 days of dosing for both active moieties. Aripiprazole accumulation is predictable from single-dose pharmacokinetics. At steady state, the pharmacokinetics of aripiprazole are dose-proportional. Elimination of aripiprazole is mainly through hepatic metabolism involving two P450 isozymes, CYP2D6 and CYP3A4.

Absorption

Aripiprazole is well absorbed, with peak plasma concentrations occurring within 3 to 5 hours; the absolute oral bioavailability of the tablet formulation is 87%. ABILIFY can be administered with or without food. Administration of a 15-mg ABILIFY tablet with a standard high-fat meal did not significantly affect the C_{max} or AUC of aripiprazole or its active metabolite, dehydro-aripiprazole, but delayed T_{max} by 3 hours for aripiprazole and 12 hours for dehydro-aripiprazole.

Distribution

The steady-state volume of distribution of aripiprazole following intravenous administration is high (404 L or 4.9 L/kg), indicating extensive extravascular distribution. At therapeutic concentrations, aripiprazole and its major metabolite are greater than 99% bound to serum proteins, primarily to albumin. In healthy human volunteers administered 0.5 to 30 mg/day aripiprazole for 14 days, there was dose-dependent D_2 -receptor occupancy indicating brain penetration of aripiprazole in humans.

Metabolism and Elimination

Aripiprazole is metabolized primarily by three biotransformation pathways: dehydrogenation, hydroxylation, and N-dealkylation. Based on *in vitro* studies, CYP3A4

and CYP2D6 enzymes are responsible for dehydrogenation and hydroxylation of aripiprazole, and N-dealkylation is catalyzed by CYP3A4. Aripiprazole is the predominant drug moiety in the systemic circulation. At steady state, dehydro-aripiprazole, the active metabolite, represents about 40% of aripiprazole AUC in plasma.

Approximately 8% of Caucasians lack the capacity to metabolize CYP2D6 substrates and are classified as poor metabolizers (PM), whereas the rest are extensive metabolizers (EM). PMs have about an 80% increase in aripiprazole exposure and about a 30% decrease in exposure to the active metabolite compared to EMs, resulting in about a 60% higher exposure to the total active moieties from a given dose of aripiprazole compared to EMs. Coadministration of ABILIFY with known inhibitors of CYP2D6, like quinidine in EMs results in a 112% increase in aripiprazole plasma exposure, and dosing adjustment is needed (see PRECAUTIONS: Drug-Drug Interactions). The mean elimination half-lives are about 75 hours and 146 hours for aripiprazole in EMs and PMs, respectively. Aripiprazole does not inhibit or induce the CYP2D6 pathway.

Following a single oral dose of [¹⁴C]-labeled aripiprazole, approximately 25% and 55% of the administered radioactivity was recovered in the urine and feces, respectively. Less than 1% of unchanged aripiprazole was excreted in the urine and approximately 18% of the oral dose was recovered unchanged in the feces.

Special Populations

In general, no dosage adjustment for ABILIFY is required on the basis of a patient's age, gender, race, smoking status, hepatic function, or renal function (see DOSAGE AND ADMINISTRATION: Dosage in Special Populations). The pharmacokinetics of aripiprazole in special populations are described below.

Hepatic Impairment

In a single-dose study (15 mg of aripiprazole) in subjects with varying degrees of liver cirrhosis (Child-Pugh Classes A, B, and C), the AUC of aripiprazole, compared to healthy subjects, increased 31% in mild HI, increased 8% in moderate HI, and decreased 20% in severe HI. None of these differences would require dose adjustment.

Renal Impairment

In patients with severe renal impairment (creatinine clearance < 30 mL/min), C_{max} of aripiprazole (given in a single dose of 15 mg) and dehydro-aripiprazole increased by 36% and 53%, respectively, but AUC was 15% lower for aripiprazole and 7% higher for dehydro-aripiprazole. Renal excretion of both unchanged aripiprazole and dehydro-aripiprazole is less than 1% of the dose. No dosage adjustment is required in subjects with renal impairment.

Elderly

In formal single-dose pharmacokinetic studies (with aripiprazole given in a single dose of 15 mg), aripiprazole clearance was 20% lower in elderly (≥ 65 years) subjects compared to younger adult subjects (18-64 years). There was no detectable age effect, however, in the population pharmacokinetic analysis in schizophrenia patients. Also, the pharmacokinetics of aripiprazole after multiple doses in elderly patients appeared similar to that observed in young healthy subjects. No dosage adjustment is recommended for elderly patients. (see PRECAUTIONS: Geriatric Use).

Gender

C_{max} and AUC of aripiprazole and its active metabolite, dehydro-aripiprazole, are 30-40% higher in women than in men, and correspondingly, the apparent oral clearance of aripiprazole is lower in women. These differences, however, are largely explained by differences in body weight (25%) between men and women. No dosage adjustment is recommended based on gender.

Race

Although no specific pharmacokinetic study was conducted to investigate the effects of race on the disposition of aripiprazole, population pharmacokinetic evaluation revealed no evidence of clinically significant race-related differences in the pharmacokinetics of aripiprazole. No dosage adjustment is recommended based on race.

Smoking

Based on studies utilizing human liver enzymes *in vitro*, aripiprazole is not a substrate for CYP1A2 and also does not undergo direct glucuronidation. Smoking should, therefore, not have an effect on the pharmacokinetics of aripiprazole. Consistent with these *in vitro* results, population pharmacokinetic evaluation did not reveal any significant pharmacokinetic differences between smokers and nonsmokers. No dosage adjustment is recommended based on smoking status.

Drug-Drug Interactions

Potential for Other Drugs to Affect ABILIFY

Aripiprazole is not a substrate of CYP1A1, CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, or CYP2E1 enzymes. Aripiprazole also does not undergo direct glucuronidation. This suggests that an interaction of aripiprazole with inhibitors or inducers of these enzymes, or other factors, like smoking, is unlikely.

Both CYP3A4 and CYP2D6 are responsible for aripiprazole metabolism. Agents that induce CYP3A4 (eg, carbamazepine) could cause an increase in aripiprazole clearance and lower blood levels. Inhibitors of CYP3A4 (eg, ketoconazole) or CYP2D6 (eg, quinidine, fluoxetine, or paroxetine) can inhibit aripiprazole elimination and cause increased blood levels.

Potential for ABILIFY to Affect Other Drugs

Aripiprazole is unlikely to cause clinically important pharmacokinetic interactions with drugs metabolized by cytochrome P450 enzymes. In *in vivo* studies, 10- to 30-mg/day doses of aripiprazole had no significant effect on metabolism by CYP2D6 (dextromethorphan), CYP2C9 (warfarin), CYP2C19 (omeprazole, warfarin), and CYP3A4 (dextromethorphan) substrates. Additionally, aripiprazole and dehydro-aripiprazole did not show potential for altering CYP1A2-mediated metabolism *in vitro* (see **Precautions: Drug-Drug Interactions**).

Aripiprazole had no clinically important interactions with the following drugs:

Famotidine: Coadministration of aripiprazole (given in a single dose of 15 mg) with a 40-mg single dose of the H₂ antagonist famotidine, a potent gastric acid blocker, decreased the solubility of aripiprazole and, hence, its rate of absorption, reducing by 37% and 21% the C_{max} of aripiprazole and dehydro-aripiprazole, respectively, and by 13% and 15%, respectively, the extent of absorption (AUC). No dosage adjustment of aripiprazole is required when administered concomitantly with famotidine.

Valproate: When valproate (500-1500 mg/day) and aripiprazole (30 mg/day) were co-administered at steady state, the C_{max} and AUC of aripiprazole were decreased by 25%. No dosage adjustment of aripiprazole is required when administered concomitantly with valproate.

Lithium: A pharmacokinetic interaction of aripiprazole with lithium is unlikely because lithium is not bound to plasma proteins, is not metabolized, and is almost entirely excreted unchanged in urine. Coadministration of therapeutic doses of lithium (1200-1800 mg/day) for 21 days with aripiprazole (30 mg/day) did not result in clinically significant changes in the pharmacokinetics of aripiprazole or its active metabolite, dehydro-aripiprazole (C_{max} and AUC increased by less than 20%). No dosage adjustment of aripiprazole is required when administered concomitantly with lithium.

Dextromethorphan: Aripiprazole at doses of 10 to 30 mg per day for 14 days had no effect on dextromethorphan's O-dealkylation to its major metabolite, dextrorphan, a pathway known to be dependent on CYP2D6 activity. Aripiprazole also had no effect on dextromethorphan's N-demethylation to its metabolite 3-methoxymorphan, a pathway known to be dependent on CYP3A4 activity. No dosage adjustment of dextromethorphan is required when administered concomitantly with aripiprazole.

Warfarin: Aripiprazole 10 mg per day for 14 days had no effect on the pharmacokinetics of R- and S-warfarin or on the pharmacodynamic end point of International Normalized Ratio, indicating the lack of a clinically relevant effect of aripiprazole on CYP2C9 and CYP2C19 metabolism or the binding of highly protein-bound warfarin. No dosage adjustment of warfarin is required when administered concomitantly with aripiprazole.

Omeprazole: Aripiprazole 10 mg per day for 15 days had no effect on the pharmacokinetics of a single 20-mg dose of omeprazole, a CYP2C19 substrate, in healthy subjects. No dosage adjustment of omeprazole is required when administered concomitantly with aripiprazole.

Clinical Studies

The efficacy of ABILIFY in the treatment of schizophrenia was evaluated in four short-term (4- and 6-week), placebo-controlled trials of acutely relapsed inpatients who predominantly met DSM-III/IV criteria for schizophrenia. Three of the four trials were able to distinguish aripiprazole from placebo, but one study, the smallest, did not. Three of these studies also included an active control group consisting of either risperidone (one trial) or haloperidol (two trials), but they were not designed to allow for a comparison of ABILIFY and the active comparators.

In the three positive trials for ABILIFY, four primary measures were used for assessing psychiatric signs and symptoms. The Positive and Negative Syndrome Scale (PANSS) is a multi-item inventory of general psychopathology used to evaluate the effects of drug treatment in schizophrenia. The PANSS positive subscale is a subset of items in the PANSS that rates seven positive symptoms of schizophrenia (delusions, conceptual disorganization, hallucinatory behavior, excitement, grandiosity, suspiciousness/persecution, and hostility). The PANSS negative subscale is a subset of items in the PANSS that rates seven negative symptoms of schizophrenia (blunted affect, emotional withdrawal, poor rapport, passive apathetic withdrawal, difficulty in abstract thinking, lack of spontaneity/flow of conversation, stereotyped thinking). The Clinical Global Impression (CGI) assessment reflects the impression of a skilled observer, fully familiar with the manifestations of schizophrenia, about the overall clinical state of the patient.

In a 4-week trial (n=414) comparing two fixed doses of ABILIFY (15 or 30 mg/day) and haloperidol (10 mg/day) to placebo, both doses of ABILIFY were superior to placebo in the PANSS total score, PANSS positive subscale, and CGI-severity score. In addition, the 15-mg dose was superior to placebo in the PANSS negative subscale.

In a 4-week trial (n=404) comparing two fixed doses of ABILIFY (20 or 30 mg/day) and risperidone (6 mg/day) to placebo, both doses of ABILIFY were superior to placebo in the PANSS total score, PANSS positive subscale, PANSS negative subscale, and CGI-severity score.

In a 6-week trial (n=420) comparing three fixed doses of ABILIFY (10, 15 or 20 mg/day) to placebo, all three doses of ABILIFY were superior to placebo in the PANSS total score, PANSS positive subscale, and the PANSS negative subscale.

In a fourth study, a 4-week trial (n=103) comparing ABILIFY in a range of 5 to 30 mg/day or haloperidol 5 to 20 mg/day to placebo, haloperidol was superior to placebo, in the Brief Psychiatric Rating Scale (BPRS), a multi-item inventory of general

psychopathology traditionally used to evaluate the effects of drug treatment in psychosis, and in a responder analysis based on the CGI-severity score, the primary outcomes for that trial. ABILIFY was only significantly different compared to placebo in a responder analysis based on the CGI-severity score.

Thus, the efficacy of 15-mg, 20-mg, and 30-mg daily doses was established in two studies for each dose whereas the efficacy of the 10-mg dose was established in one study. There was no evidence in any study that the higher dose groups offered any advantage over the lowest dose group.

An examination of population subgroups did not reveal any clear evidence of differential responsiveness on the basis of age, gender, or race.

INDICATIONS AND USAGE

ABILIFY is indicated for the treatment of schizophrenia. The efficacy of ABILIFY in the treatment of schizophrenia was established in short-term (4- and 6-week) controlled trials of schizophrenic inpatients (see **CLINICAL PHARMACOLOGY: Clinical Studies**).

The long-term efficacy of aripiprazole in the treatment of schizophrenia has not been established. The physician who elects to use ABILIFY for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient.

CONTRAINDICATIONS

ABILIFY is contraindicated in patients with a known hypersensitivity to the product.

WARNINGS

Neuroleptic Malignant Syndrome (NMS)

A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with administration of antipsychotic drugs, including aripiprazole. Two possible cases of NMS occurred during aripiprazole treatment in the premarketing worldwide clinical database. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure.

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to exclude cases where the clinical presentation includes both serious medical illness (eg, pneumonia, systemic infection, etc.) and untreated or

inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system pathology.

The management of NMS should include: 1) immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy; 2) intensive symptomatic treatment and medical monitoring; and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for uncomplicated NMS.

If a patient requires antipsychotic drug treatment after recovery from NMS, the potential reintroduction of drug therapy should be carefully considered. The patient should be carefully monitored, since recurrences of NMS have been reported.

Tardive Dyskinesia

A syndrome of potentially irreversible, involuntary, dyskinetic movements may develop in patients treated with antipsychotic drugs. Although the prevalence of the syndrome appears to be highest among the elderly, especially elderly women, it is impossible to rely upon prevalence estimates to predict, at the inception of antipsychotic treatment, which patients are likely to develop the syndrome. Whether antipsychotic drug products differ in their potential to cause tardive dyskinesia is unknown.

The risk of developing tardive dyskinesia and the likelihood that it will become irreversible are believed to increase as the duration of treatment and the total cumulative dose of antipsychotic drugs administered to the patient increase. However, the syndrome can develop, although much less commonly, after relatively brief treatment periods at low doses.

There is no known treatment for established cases of tardive dyskinesia, although the syndrome may remit, partially or completely, if antipsychotic treatment is withdrawn. Antipsychotic treatment, itself, however, may suppress (or partially suppress) the signs and symptoms of the syndrome and, thereby, may possibly mask the underlying process. The effect that symptomatic suppression has upon the long-term course of the syndrome is unknown.

Given these considerations, ABILIFY should be prescribed in a manner that is most likely to minimize the occurrence of tardive dyskinesia. Chronic antipsychotic treatment should generally be reserved for patients who suffer from a chronic illness that (1) is known to respond to antipsychotic drugs, and (2) for whom alternative, equally effective, but potentially less harmful treatments are not available or appropriate. In patients who do require chronic treatment, the smallest dose and the shortest duration of treatment producing a satisfactory clinical response should be sought. The need for continued treatment should be reassessed periodically.

If signs and symptoms of tardive dyskinesia appear in a patient on ABILIFY, drug discontinuation should be considered. However, some patients may require treatment with ABILIFY despite the presence of the syndrome.

PRECAUTIONS

General

Orthostatic Hypotension

Aripiprazole may be associated with orthostatic hypotension, perhaps due to its α_1 -adrenergic receptor antagonism. The incidence of orthostatic hypotension associated events from five short-term, placebo-controlled trials in schizophrenia (n=926) on Abilify included: orthostatic hypotension (placebo 1%, aripiprazole 1.9%), orthostatic lightheadedness (placebo 1%, aripiprazole 0.9%), and syncope (placebo 1%, aripiprazole 0.6%). The incidence of a significant orthostatic change in blood pressure (defined as a decrease of at least 30 mmHg in systolic blood pressure when changing from a supine to standing position) for aripiprazole was not statistically different from placebo (14% among aripiprazole-treated patients and 12% among placebo-treated patients).

Aripiprazole should be used with caution in patients with known cardiovascular disease (history of myocardial infarction or ischemic heart disease, heart failure or conduction abnormalities), cerebrovascular disease, or conditions which would predispose patients to hypotension (dehydration, hypovolemia, and treatment with antihypertensive medications).

Seizure

Seizures occurred in 0.1% (1/926) of aripiprazole-treated patients in short-term, placebo-controlled trials. As with other antipsychotic drugs, aripiprazole should be used cautiously in patients with a history of seizures or with conditions that lower the seizure threshold, eg, Alzheimer's dementia. Conditions that lower the seizure threshold may be more prevalent in a population of 65 years or older.

Potential for Cognitive and Motor Impairment

In short-term, placebo-controlled trials, somnolence was reported in 11% of patients on ABILIFY compared to 8% of patients on placebo; somnolence led to discontinuation in 0.1% (1/926) of patients on ABILIFY in short-term, placebo-controlled trials. Despite the relatively modest increased incidence of somnolence compared to placebo, ABILIFY, like other antipsychotics, may have the potential to impair judgment, thinking, or motor skills. Patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that therapy with ABILIFY does not affect them adversely.

Body Temperature Regulation

Disruption of the body's ability to reduce core body temperature has been attributed to antipsychotic agents. Appropriate care is advised when prescribing aripiprazole for patients who will be experiencing conditions which may contribute to an elevation in core body temperature, eg, exercising strenuously, exposure to extreme heat, receiving concomitant medication with anticholinergic activity, or being subject to dehydration.

Dysphagia

Esophageal dysmotility and aspiration have been associated with antipsychotic drug use. Aspiration pneumonia is a common cause of morbidity and mortality in elderly patients, in particular those with advanced Alzheimer's dementia. Aripiprazole and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia (see PRECAUTIONS: Use in Patients with Concomitant Illness).

Suicide

The possibility of a suicide attempt is inherent in psychotic illnesses, and close supervision of high-risk patients should accompany drug therapy. Prescriptions for ABILIFY should be written for the smallest quantity of tablets consistent with good patient management in order to reduce the risk of overdose.

Use in Patients with Concomitant Illness

Safety Experience in Elderly Patients with Psychosis Associated with Alzheimer's Disease: In a flexible dose (2 to 15 mg/day), 10-week, placebo-controlled study of aripiprazole in elderly patients (mean age: 81.5 years; range: 56-95) with psychosis associated with Alzheimer's dementia, 4 of 105 patients (3.8%) who received ABILIFY died compared to no deaths among 102 patients who received placebo during or within 30 days after termination of the double-blind portion of the study. Three of the patients (age 92, 91, and 87 years) died following the discontinuation of ABILIFY in the double-blind phase of the study (causes of death were pneumonia, heart failure, and shock). The fourth patient (age 78 years) died following hip surgery while in the double-blind portion of the study. The treatment-emergent adverse events that were reported at an incidence of $\geq 5\%$ and having a greater incidence than placebo in this study were accidental injury, somnolence, and bronchitis. Eight percent of the ABILIFY treated patients reported somnolence compared to 1% of placebo patients. In a small pilot, open-label, ascending-dose, cohort study (n=30) in elderly patients with dementia, ABILIFY was associated in a dose-related fashion with somnolence.

The safety and efficacy of ABILIFY in the treatment of patients with psychosis associated with dementia has not been established. If the prescriber elects to treat such patients with ABILIFY, vigilance should be exercised, particularly for the emergence of difficulty swallowing or excessive somnolence, which could predispose to accidental injury or aspiration.

Clinical experience with ABILIFY in patients with certain concomitant systemic illnesses (see **CLINICAL PHARMACOLOGY: Special Populations: Renal Impairment and Hepatic Impairment**) is limited.

ABILIFY has not been evaluated or used to any appreciable extent in patients with a recent history of myocardial infarction or unstable heart disease. Patients with these diagnoses were excluded from premarketing clinical studies.

Information for Patients

Physicians are advised to discuss the following issues with patients for whom they prescribe ABILIFY:

Interference with Cognitive and Motor Performance

Because aripiprazole may have the potential to impair judgment, thinking, or motor skills, patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that aripiprazole therapy does not affect them adversely.

Pregnancy

Patients should be advised to notify their physician if they become pregnant or intend to become pregnant during therapy with ABILIFY.

Nursing

Patients should be advised not to breast-feed an infant if they are taking ABILIFY.

Concomitant Medication

Patients should be advised to inform their physicians if they are taking, or plan to take, any prescription or over-the-counter drugs, since there is a potential for interactions.

Alcohol

Patients should be advised to avoid alcohol while taking ABILIFY.

Heat Exposure and Dehydration

Patients should be advised regarding appropriate care in avoiding overheating and dehydration.

Drug-Drug Interactions

Given the primary CNS effects of aripiprazole, caution should be used when ABILIFY is taken in combination with other centrally acting drugs and alcohol. Due to its α_1 -adrenergic receptor antagonism, aripiprazole has the potential to enhance the effect of certain antihypertensive agents.

Potential for Other Drugs to Affect ABILIFY

Aripiprazole is not a substrate of CYP1A1, CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, or CYP2E1 enzymes. Aripiprazole also does not undergo direct glucuronidation. This suggests that an interaction of aripiprazole with inhibitors or inducers of these enzymes, or other factors, like smoking, is unlikely.

Both CYP3A4 and CYP2D6 are responsible for aripiprazole metabolism. Agents that induce CYP3A4 (eg, carbamazepine) could cause an increase in aripiprazole clearance and lower blood levels. Inhibitors of CYP3A4 (eg, ketoconazole) or CYP2D6 (eg, quinidine, fluoxetine, or paroxetine) can inhibit aripiprazole elimination and cause increased blood levels.

Ketoconazole: Coadministration of ketoconazole (200 mg/day for 14 days) with a 15-mg single dose of aripiprazole increased the AUC of aripiprazole and its active metabolite by 63% and 77%, respectively. The effect of a higher ketoconazole dose (400 mg/day) has not been studied. When concomitant administration of ketoconazole with aripiprazole occurs, aripiprazole dose should be reduced to one-half of its normal dose. Other strong inhibitors of CYP3A4 (itraconazole) would be expected to have similar effects and need similar dose reductions; weaker inhibitors (erythromycin, grapefruit juice) have not been studied. When the CYP3A4 inhibitor is withdrawn from the combination therapy, aripiprazole dose should then be increased.

Quinidine: Coadministration of a 10-mg single dose of aripiprazole with quinidine (166 mg/day for 13 days), a potent inhibitor of CYP2D6, increased the AUC of aripiprazole by 112% but decreased the AUC of its active metabolite, dehydro-aripiprazole, by 35%. Aripiprazole dose should be reduced to one-half of its normal dose when concomitant administration of quinidine with aripiprazole occurs. Other significant inhibitors of CYP2D6, such as fluoxetine or paroxetine, would be expected to have similar effects and, therefore, should be accompanied by similar dose reductions. When the CYP2D6 inhibitor is withdrawn from the combination therapy, aripiprazole dose should then be increased.

Carbamazepine: Coadministration of carbamazepine (200 mg BID), a potent CYP3A4 inducer, with aripiprazole (30 mg QD) resulted in an approximate 70% decrease in C_{max} and AUC values of both aripiprazole and its active metabolite, dehydro-aripiprazole. When carbamazepine is added to aripiprazole therapy, aripiprazole dose should be doubled. Additional dose increases should be based on clinical evaluation. When carbamazepine is withdrawn from the combination therapy, aripiprazole dose should then be reduced.

No clinically significant effect of famotidine, valproate, or lithium was seen on the pharmacokinetics of aripiprazole (see **CLINICAL PHARMACOLOGY: Drug-Drug Interactions**).

Potential for ABILIFY to Affect Other Drugs

Aripiprazole is unlikely to cause clinically important pharmacokinetic interactions with drugs metabolized by cytochrome P450 enzymes. In *in vivo* studies, 10- to 30-mg/day doses of aripiprazole had no significant effect on metabolism by CYP2D6 (dextromethorphan), CYP2C9 (warfarin), CYP2C19 (omeprazole, warfarin), and CYP3A4 (dextromethorphan) substrates. Additionally, aripiprazole and dehydro-aripiprazole did not show potential for altering CYP1A2-mediated metabolism *in vitro* (see **Clinical Pharmacology: Drug-Drug Interactions**).

Alcohol: There was no significant difference between aripiprazole coadministered with ethanol and placebo coadministered with ethanol on performance of gross motor skills or stimulus response in healthy subjects. As with most psychoactive medications, patients should be advised to avoid alcohol while taking ABILIFY.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Lifetime carcinogenicity studies were conducted in ICR mice and in Sprague-Dawley (SD) and F344 rats. Aripiprazole was administered for 2 years in the diet at doses of 1, 3, 10, and 30 mg/kg/day to ICR mice and 1, 3, and 10 mg/kg/day to F344 rats (0.2 to 5 and 0.3 to 3 times the maximum recommended human dose [MRHD] based on mg/m², respectively). In addition, SD rats were dosed orally for 2 years at 10, 20, 40, and 60 mg/kg/day (3 to 19 times the MRHD based on mg/m²). Aripiprazole did not induce tumors in male mice or rats. In female mice, the incidences of pituitary gland adenomas and mammary gland adenocarcinomas and adenoacanthomas were increased at dietary doses of 3 to 30 mg/kg/day (0.1 to 0.9 times human exposure at MRHD based on AUC and 0.5 to 5 times the MRHD based on mg/m²). In female rats, the incidence of mammary gland fibroadenomas was increased at a dietary dose of 10 mg/kg/day (0.1 times human exposure at MRHD based on AUC and 3 times the MRHD based on mg/m²); and the incidences of adrenocortical carcinomas and combined adrenocortical adenomas/carcinomas were increased at an oral dose of 60 mg/kg/day (14 times human exposure at MRHD based on AUC and 19 times the MRHD based on mg/m²).

Proliferative changes in the pituitary and mammary gland of rodents have been observed following chronic administration of other antipsychotic agents and are considered prolactin-mediated. Serum prolactin was not measured in the aripiprazole carcinogenicity studies. However, increases in serum prolactin levels were observed in female mice in a 13-week dietary study at the doses associated with mammary gland and pituitary tumors. Serum prolactin was not increased in female rats in 4- and 13-week

dietary studies at the dose associated with mammary gland tumors. The relevance for human risk of the findings of prolactin-mediated endocrine tumors in rodents is unknown.

Mutagenesis

The mutagenic potential of aripiprazole was tested in the *in vitro* bacterial reverse-mutation assay, the *in vitro* bacterial DNA repair assay, the *in vitro* forward gene mutation assay in mouse lymphoma cells, the *in vitro* chromosomal aberration assay in Chinese hamster lung (CHL) cells, the *in vivo* micronucleus assay in mice, and the unscheduled DNA synthesis assay in rats. Aripiprazole and a metabolite (2,3-DCPP) were clastogenic in the *in vitro* chromosomal aberration assay in CHL cells with and without metabolic activation. The metabolite, 2,3-DCPP, produced increases in numerical aberrations in the *in vitro* assay in CHL cells in the absence of metabolic activation. A positive response was obtained in the *in vivo* micronucleus assay in mice, however, the response was shown to be due to a mechanism not considered relevant to humans.

Impairment of Fertility

Female rats were treated with oral doses of 2, 6, and 20 mg/kg/day (0.6, 2, and 6 times the maximum recommended human dose [MRHD] on a mg/m² basis) of aripiprazole from 2 weeks prior to mating through day 7 of gestation. Estrus cycle irregularities and increased corpora lutea were seen at all doses, but no impairment of fertility was seen. Increased pre-implantation loss was seen at 6 and 20 mg/kg, and decreased fetal weight was seen at 20 mg/kg.

Male rats were treated with oral doses of 20, 40, and 60 mg/kg/day (6, 13, and 19 times the MRHD on a mg/m² basis) of aripiprazole from 9 weeks prior to mating through mating. Disturbances in spermatogenesis were seen at 60 mg/kg, and prostate atrophy was seen at 40 and 60 mg/kg, but no impairment of fertility was seen.

Pregnancy

Pregnancy Category C

In animal studies aripiprazole demonstrated developmental toxicity, including possible teratogenic effects in rats and rabbits.

Pregnant rats were treated with oral doses of 3, 10, and 30 mg/kg/day (1, 3, and 10 times the maximum recommended human dose [MRHD] on a mg/m² basis) of aripiprazole during the period of organogenesis. Gestation was slightly prolonged at 30 mg/kg. Treatment caused a slight delay in fetal development as evidenced by decreased fetal weight (30 mg/kg), undescended testes (30 mg/kg), and delayed skeletal ossification (10 and 30 mg/kg). There were no adverse effects on embryofetal or pup survival. Delivered offspring had decreased bodyweights (10 and 30 mg/kg), and increased incidences of hepatodiaphragmatic nodules and diaphragmatic hernia at 30 mg/kg (the other dose groups were not examined for these findings). (A low incidence of diaphragmatic hernia

was also seen in the fetuses exposed to 30 mg/kg.) Postnatally, delayed vaginal opening was seen at 10 and 30 mg/kg and impaired reproductive performance (decreased fertility rate, corpora lutea, implants, and live fetuses, and increased post-implantation loss, likely mediated through effects on female offspring) was seen at 30 mg/kg. Some maternal toxicity was seen at 30 mg/kg, however, there was no evidence to suggest that these developmental effects were secondary to maternal toxicity.

Pregnant rabbits were treated with oral doses of 10, 30, and 100 mg/kg/day (2, 3, and 11 times human exposure at MRHD based on AUC and 6, 19, and 65 times the MRHD based on mg/m^2) of aripiprazole during the period of organogenesis. Decreased maternal food consumption and increased abortions were seen at 100 mg/kg. Treatment caused increased fetal mortality (100 mg/kg), decreased fetal weight (30 and 100 mg/kg), increased incidence of a skeletal abnormality (fused sternbrae at 30 and 100 mg/kg) and minor skeletal variations (100 mg/kg).

In a study in which rats were treated with oral doses of 3, 10, and 30 mg/kg/day (1, 3, and 10 times the MRHD on a mg/m^2 basis) of aripiprazole perinatally and postnatally (from day 17 of gestation through day 21 postpartum), slight maternal toxicity and slightly prolonged gestation were seen at 30 mg/kg. An increase in stillbirths, and decreases in pup weight (persisting into adulthood) and survival, were seen at this dose.

There are no adequate and well-controlled studies in pregnant women. It is not known whether aripiprazole can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. Aripiprazole should be used during pregnancy only if the potential benefit outweighs the potential risk to the fetus.

Labor and Delivery

The effect of aripiprazole on labor and delivery in humans is unknown.

Nursing Mothers

Aripiprazole was excreted in milk of rats during lactation. It is not known whether aripiprazole or its metabolites are excreted in human milk. It is recommended that women receiving aripiprazole should not breast-feed.

Pediatric Use

Safety and effectiveness in pediatric and adolescent patients have not been established.

Geriatric Use

Of the 5592 patients treated with aripiprazole in premarketing clinical trials, 659 (12%) were ≥ 65 years old and 525 (9%) were ≥ 75 years old. The majority (91%) of the 659 patients were diagnosed with dementia of the Alzheimer's type.

Placebo-controlled studies of aripiprazole in schizophrenia did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. There was no effect of age on the pharmacokinetics of a single 15-mg dose of aripiprazole. Aripiprazole clearance was decreased by 20% in elderly subjects (≥ 65 years) compared to younger adult subjects (18-64 years), but there was no detectable effect of age in the population pharmacokinetic analysis in schizophrenia patients.

Studies of elderly patients with psychosis associated with Alzheimer's disease have suggested that there may be a different tolerability profile in this population compared to younger patients with schizophrenia (see **PRECAUTIONS: Use in Patients with Concomitant Illness**). The safety and efficacy of ABILIFY in the treatment of patients with psychosis associated with Alzheimer's disease has not been established. If the prescriber elects to treat such patients with ABILIFY, vigilance should be exercised.

ADVERSE REACTIONS

Aripiprazole has been evaluated for safety in 5592 patients who participated in multiple-dose, premarketing trials in schizophrenia, bipolar mania, and dementia of the Alzheimer's type, and who had approximately 3639 patient-years of exposure. A total of 1887 aripiprazole-treated patients were treated for at least 180 days and 1251 aripiprazole-treated patients had at least 1 year of exposure.

The conditions and duration of treatment with aripiprazole included (in overlapping categories) double-blind and comparative and noncomparative open-label studies, inpatient and outpatient studies, fixed- and flexible-dose studies, and short- and longer-term exposure.

Adverse events during exposure were obtained by collecting volunteered adverse events, as well as results of physical examinations, vital signs, weights, laboratory analyses, and ECG. Adverse experiences were recorded by clinical investigators using terminology of their own choosing. In the tables and tabulations that follow, modified COSTART dictionary terminology has been used initially to classify reported adverse events into a smaller number of standardized event categories, in order to provide a meaningful estimate of the proportion of individuals experiencing adverse events.

The stated frequencies of adverse events represent the proportion of individuals who experienced at least once, a treatment-emergent adverse event of the type listed. An event was considered treatment emergent if it occurred for the first time or worsened while receiving therapy following baseline evaluation. There was no attempt to use investigator causality assessments; i.e., all reported events are included.

The prescriber should be aware that the figures in the tables and tabulations cannot be used to predict the incidence of side effects in the course of usual medical practice where patient characteristics and other factors differ from those that prevailed in the clinical trials. Similarly, the cited frequencies cannot be compared with figures obtained from

other clinical investigations involving different treatment, uses, and investigators. The cited figures, however, do provide the prescribing physician with some basis for estimating the relative contribution of drug and nondrug factors to the adverse event incidence in the population studied.

Adverse Findings Observed in Short-Term, Placebo-Controlled Trials of Patients with Schizophrenia

The following findings are based on a pool of five placebo-controlled trials (four 4-week and one 6-week) in which aripiprazole was administered in doses ranging from 2 to 30 mg/day.

Adverse Events Associated with Discontinuation of Treatment in Short-Term, Placebo-Controlled Trials

Overall, there was no difference in the incidence of discontinuation due to adverse events between aripiprazole-treated (7%) and placebo-treated (9%) patients. The types of adverse events that led to discontinuation were similar between the aripiprazole and placebo-treated patients.

Adverse Events Occurring at an Incidence of 2% or More Among Aripiprazole-Treated Patients and Greater than Placebo in Short-Term Placebo-Controlled Trials

Table 1 enumerates the incidence, rounded to the nearest percent, of treatment-emergent adverse events that occurred during acute therapy (up to 6 weeks), including only those events that occurred in 2% or more of patients treated with aripiprazole (doses \geq 2 mg/day) and for which the incidence in patients treated with aripiprazole was greater than the incidence in patients treated with placebo.

**Table 1: Treatment-Emergent Adverse Events
in Short-Term, Placebo-Controlled Trials**

Body System Adverse Event	Percentage of Patients Reporting Event ^a	
	Aripiprazole (n=926)	Placebo (n=413)
Body as a Whole		
Headache	32	25
Asthenia	7	5
Fever	2	1
Digestive System		
Nausea	14	10
Vomiting	12	7
Constipation	10	8
Nervous System		
Anxiety	25	24
Insomnia	24	19
Lightheadedness	11	7
Somnolence	11	8
Akathisia	10	7
Tremor	3	2
Respiratory System		
Rhinitis	4	3
Coughing	3	2
Skin and Appendages		
Rash	6	5
Special Senses		
Blurred vision	3	1

^a Events reported by at least 2% of patients treated with aripiprazole, except the following events, which had an incidence equal to or less than placebo: abdominal pain, accidental injury, back pain, dental pain, dyspepsia, diarrhea, dry mouth, myalgia, agitation, psychosis, extrapyramidal syndrome, hypertonia, pharyngitis, upper respiratory tract infection, dysmenorrhea, vaginitis.

An examination of population subgroups did not reveal any clear evidence of differential adverse event incidence on the basis of age, gender, or race.

Dose-related Adverse Events

Dose response relationships for the incidence of treatment-emergent adverse events were evaluated from four trials comparing various fixed doses (2, 10, 15, 20, and 30 mg/day) of aripiprazole to placebo. This analysis, stratified by study, indicated that the only adverse event to have a possible dose response relationship, and then most prominent only with 30 mg, was somnolence (placebo, 7.7%; 15 mg, 8.7%; 20 mg, 7.5%; 30 mg, 15.3%).

Extrapyramidal Symptoms

In the short-term, placebo-controlled trials, the incidence of reported EPS for aripiprazole-treated patients was 6% vs. 6% for placebo. Objectively collected data from those trials on the Simpson Angus Rating Scale (for EPS), the Barnes Akathisia Scale (for akathisia), and the Assessments of Involuntary Movement Scales (for dyskinesias) also did not show a difference between aripiprazole and placebo, with the exception of the Barnes Akathisia Scale (aripiprazole, 0.08; placebo, -0.05).

Laboratory Test Abnormalities

A between group comparison for 4- to 6-week placebo-controlled trials revealed no medically important differences between the aripiprazole and placebo groups in the proportions of patients experiencing potentially clinically significant changes in routine serum chemistry, hematology, or urinalysis parameters. Similarly, there were no aripiprazole/placebo differences in the incidence of discontinuations for changes in serum chemistry, hematology, or urinalysis.

Weight Gain

In short-term trials, there was a slight difference in mean weight gain between aripiprazole and placebo patients (+0.7 kg vs. -0.05 kg, respectively), and also a difference in the proportion of patients meeting a weight gain criterion of $\geq 7\%$ of body weight [aripiprazole (8%) compared to placebo (3%)]. The following table provides the weight change results from a long-term (52-week) study of aripiprazole, both mean change from baseline and proportions of patients meeting a weight gain criterion of $\geq 7\%$ of body weight relative to baseline, categorized by BMI at baseline:

Table 2: Weight Change Results Categorized by BMI at Baseline

	BMI<23	BMI 23-27	BMI>27
Mean change from baseline (kg)	2.6	1.4	-1.2
% with $\geq 7\%$ increase BW	30%	19%	8%

ECG Changes

Between group comparisons for pooled, placebo-controlled trials revealed no significant differences between aripiprazole and placebo in the proportion of patients experiencing potentially important changes in ECG parameters; in fact, within the dose range of 10 to 30 mg/day, aripiprazole tended to slightly shorten the QTc interval. Aripiprazole was associated with a median increase in heart rate of 4 beats per minute compared to a 1 beat per minute increase among placebo patients.

Other Adverse Events Observed During the Premarketing Evaluation of Aripiprazole

Following is a list of modified COSTART terms that reflect treatment-emergent adverse events as defined in the introduction to the **ADVERSE REACTIONS** section reported by patients treated with aripiprazole at multiple doses ≥ 2 mg/day during any phase of a trial within the database of 5592 patients. All reported events are included except those already listed in Table 1, or other parts of the **ADVERSE REACTIONS** section, those considered in the **WARNINGS** or **PRECAUTIONS**, those event terms which were so general as to be uninformative, events reported with an incidence of $<0.05\%$ and which did not have a substantial probability of being acutely life-threatening, events that are otherwise common as background events, and events considered unlikely to be drug related. It is important to emphasize that, although the events reported occurred during treatment with aripiprazole, they were not necessarily caused by it.

Events are further categorized by body system and listed in order of decreasing frequency according to the following definitions: frequent adverse events are those occurring in at least 1/100 patients (only those not already listed in the tabulated results from placebo-controlled trials appear in this listing); infrequent adverse events are those occurring in 1/100 to 1/1000 patients; rare events are those occurring in fewer than 1/1000 patients.

Body as a Whole: Frequent - flu syndrome, peripheral edema, chest pain, neck pain, neck rigidity; *Infrequent* - pelvic pain, suicide attempt, face edema, malaise, photosensitivity, arm rigidity, jaw pain, chills, bloating, jaw tightness, enlarged abdomen, chest tightness; *Rare* - throat pain, back tightness, head heaviness, moniliasis, throat tightness, leg rigidity, neck tightness, Mendelson's syndrome, heat stroke.

Cardiovascular System: Frequent - hypertension, tachycardia, hypotension, bradycardia; *Infrequent* - palpitation, hemorrhage, myocardial infarction, prolonged QT interval, cardiac arrest, atrial fibrillation, heart failure, AV block, myocardial ischemia, phlebitis, deep vein thrombosis, angina pectoris, extrasystoles; *Rare* - vasovagal reaction, cardiomegaly, atrial flutter, thrombophlebitis.

Digestive System: Frequent - anorexia, nausea and vomiting; *Infrequent* - increased appetite, gastroenteritis, dysphagia, flatulence, gastritis, tooth caries, gingivitis, hemorrhoids, gastroesophageal reflux, gastrointestinal hemorrhage, periodontal abscess, tongue edema, fecal incontinence, colitis, rectal hemorrhage, stomatitis, mouth ulcer, cholecystitis, fecal impaction, oral moniliasis, cholelithiasis, eructation, intestinal obstruction, peptic ulcer; *Rare* - esophagitis, gum hemorrhage, glossitis, hematemesis, melena, duodenal ulcer, cheilitis, hepatitis, hepatomegaly, pancreatitis, intestinal perforation.

Endocrine System: Infrequent - hypothyroidism; *Rare* - goiter, hyperthyroidism.

Hemic/Lymphatic System: Frequent - ecchymosis, anemia; *Infrequent* - hypochromic anemia, leukopenia, leukocytosis, lymphadenopathy, thrombocytopenia; *Rare* - eosinophilia, thrombocythemia, macrocytic anemia.

Metabolic and Nutritional Disorders: Frequent - weight loss, creatine phosphokinase increased; *Infrequent* - dehydration, edema, hypercholesteremia, hyperglycemia, hypokalemia, diabetes mellitus, SGPT increased, hyperlipemia, hypoglycemia, thirst, BUN increased, hyponatremia, SGOT increased, alkaline phosphatase increased, iron deficiency anemia, creatinine increased, bilirubinemia, lactic dehydrogenase increased, obesity; *Rare* - hyperkalemia, gout, hypernatremia, cyanosis, hyperuricemia, hypoglycemic reaction.

Musculoskeletal System: Frequent - muscle cramp; *Infrequent* - arthralgia, bone pain, myasthenia, arthritis, arthrosis, muscle weakness, spasm, bursitis; *Rare* - rhabdomyolysis, tendonitis, tenosynovitis, rheumatoid arthritis, myopathy.

Nervous System: Frequent - depression, nervousness, increased salivation, hostility, suicidal thought, manic reaction, abnormal gait, confusion, cogwheel rigidity; *Infrequent* - dystonia, twitch, impaired concentration, paresthesia, vasodilation, hypesthesia, extremity tremor, impotence, bradykinesia, decreased libido, panic attack, apathy, dyskinesia, hypersomnia, vertigo, dysarthria, tardive dyskinesia, ataxia, impaired memory, stupor, increased libido, amnesia, cerebrovascular accident, hyperactivity, depersonalization, hypokinesia, restless leg, myoclonus, dysphoria, neuropathy, increased reflexes, slowed thinking, hyperkinesia, hyperesthesia, hypotonia, oculogyric crisis; *Rare* - delirium, euphoria, buccoglossal syndrome, akinesia, blunted affect, decreased consciousness, incoordination, cerebral ischemia, decreased reflexes, obsessive thought, intracranial hemorrhage.

Respiratory System: Frequent - dyspnea, pneumonia; *Infrequent* - asthma, epistaxis, hiccup, laryngitis; *Rare* - hemoptysis, aspiration pneumonia, increased sputum, dry nasal passages, pulmonary edema, pulmonary embolism, hypoxia, respiratory failure, apnea.

Skin and Appendages: Frequent - dry skin, pruritus, sweating, skin ulcer; *Infrequent* - acne, vesiculobullous rash, eczema, alopecia, psoriasis, seborrhea; *Rare* - maculopapular rash, exfoliative dermatitis, urticaria.

Special Senses: Frequent - conjunctivitis, ear pain; *Infrequent* - dry eye, eye pain, tinnitus, otitis media, cataract, altered taste, blepharitis; *Rare* - increased lacrimation, frequent blinking, otitis externa, amblyopia, deafness, diplopia, eye hemorrhage, photophobia.

Urogenital System: Frequent - urinary incontinence; *Infrequent* - cystitis, urinary frequency, leukorrhea, urinary retention, hematuria, dysuria, amenorrhea, abnormal ejaculation, vaginal hemorrhage, vaginal moniliasis, kidney failure, uterus hemorrhage, menorrhagia, albuminuria, kidney calculus, nocturia, polyuria, urinary urgency; *Rare* -

breast pain, cervicitis, female lactation, anorgasmia, urinary burning, glycosuria, gynecomastia, urolithiasis, priapism.

DRUG ABUSE AND DEPENDENCE

Controlled Substance

ABILIFY (aripiprazole) is not a controlled substance.

Abuse and Dependence

Aripiprazole has not been systematically studied in humans for its potential for abuse, tolerance, or physical dependence. In physical dependence studies in monkeys, withdrawal symptoms were observed upon abrupt cessation of dosing. While the clinical trials did not reveal any tendency for any drug-seeking behavior, these observations were not systematic and it is not possible to predict on the basis of this limited experience the extent to which a CNS-active drug will be misused, diverted, and/or abused once marketed. Consequently, patients should be evaluated carefully for a history of drug abuse, and such patients should be observed closely for signs of ABILIFY misuse or abuse (eg, development of tolerance, increases in dose, drug-seeking behavior).

OVERDOSAGE

Human Experience

In premarketing clinical studies, involving more than 5500 patients, accidental or intentional acute overdosage of aripiprazole was identified in seven patients. In the two patients taking the largest identified amount, 180 mg, the only symptoms reported were somnolence and vomiting in one of the two patients. In the patients who were evaluated in hospital settings, including the two patients taking 180 mg, there were no observations indicating an adverse change in vital signs, laboratory assessments or ECG. An uneventful, accidental overdose (15 mg) occurred in a non-patient, an 18-month-old child, with concomitant ingestion of Ativan[®] (2 mg).

Management of Overdosage

No specific information is available on the treatment of overdose with aripiprazole. An electrocardiogram should be obtained in case of overdosage and, if QT_c interval prolongation is present, cardiac monitoring should be instituted. Otherwise, management of overdose should concentrate on supportive therapy, maintaining an adequate airway, oxygenation and ventilation, and management of symptoms. Close medical supervision and monitoring should continue until the patient recovers.

ATIVAN[®] is a registered trademark of Wyeth Laboratories, a Wyeth-Ayerst Company.

Charcoal: In the event of an overdose of ABILIFY, an early charcoal administration may be useful in partially preventing the absorption of aripiprazole. Administration of 50 g of activated charcoal, one hour after a single 15-mg oral dose of aripiprazole, decreased the mean AUC and Cmax of aripiprazole by 50%.

Hemodialysis: Although there is no information on the effect of hemodialysis in treating an overdose with aripiprazole, hemodialysis is unlikely to be useful in overdose management since aripiprazole is highly bound to plasma proteins.

DOSAGE AND ADMINISTRATION

Usual Dose

The recommended starting and target dose for ABILIFY is 10 or 15 mg/day administered on a once-a-day schedule without regard to meals. ABILIFY has been systematically evaluated and shown to be effective in a dose range of 10 to 30 mg/day, however, doses higher than 10 or 15 mg/day, the lowest doses in these trials, were not more effective than 10 or 15 mg/day. Dosage increases should not be made before 2 weeks, the time needed to achieve steady state.

Dosage in Special Populations

Dosage adjustments are not routinely indicated on the basis of age, gender, race, or renal or hepatic impairment status (see **CLINICAL PHARMACOLOGY: Special Populations**).

Dosage adjustment for patients taking aripiprazole concomitantly with potential CYP3A4 inhibitors: When concomitant administration of ketoconazole with aripiprazole occurs, aripiprazole dose should be reduced to one-half of the usual dose. When the CYP3A4 inhibitor is withdrawn from the combination therapy, aripiprazole dose should then be increased.

Dosage adjustment for patients taking aripiprazole concomitantly with potential CYP2D6 inhibitors: When concomitant administration of potential CYP2D6 inhibitors such as quinidine, fluoxetine, or paroxetine with aripiprazole occurs, aripiprazole dose should be reduced at least to one-half of its normal dose. When the CYP2D6 inhibitor is withdrawn from the combination therapy, aripiprazole dose should then be increased.

Dosage adjustment for patients taking potential CYP3A4 inducers: When potential CYP3A4 inducer such as carbamazepine is added to aripiprazole therapy, aripiprazole dose should be doubled (to 20 or 30 mg). Additional dose increases should be based on clinical evaluation. When carbamazepine is withdrawn from the combination therapy, aripiprazole dose should be reduced to 10 to 15 mg.

Maintenance Therapy

There is no body of evidence available from controlled trials to answer the question of how long a patient treated with aripiprazole should remain on it. It is generally agreed, however, that pharmacological treatment for episodes of acute schizophrenia should continue for up to 6 months or longer. Patients should be periodically reassessed to determine the need for maintenance treatment.

Switching from Other Antipsychotics

There are no systematically collected data to specifically address switching patients with schizophrenia from other antipsychotics to ABILIFY or concerning concomitant administration with other antipsychotics. While immediate discontinuation of the previous antipsychotic treatment may be acceptable for some patients with schizophrenia, more gradual discontinuation may be most appropriate for others. In all cases, the period of overlapping antipsychotic administration should be minimized.

ANIMAL TOXICOLOGY

Aripiprazole produced retinal degeneration in albino rats in a 26-week chronic toxicity study at a dose of 60 mg/kg and in a 2-year carcinogenicity study at doses of 40 and 60 mg/kg. The 40- and 60-mg/kg doses are 13 and 19 times the maximum recommended human dose (MRHD) based on mg/m² and 7 to 14 times human exposure at MRHD based on AUC. Evaluation of the retinas of albino mice and of monkeys did not reveal evidence of retinal degeneration. Additional studies to further evaluate the mechanism have not been performed. The relevance of this finding to human risk is unknown.

HOW SUPPLIED

ABILIFY™ (aripiprazole) Tablets are available in the following strengths and packages.

The 2-mg ABILIFY tablets are green, modified rectangular, scored tablets, debossed on one side with “A-006” and “2”.

Bottles of 30	NDC 59148-006-13
Blister of 100	NDC 59148-006-35

The 5-mg ABILIFY tablets are blue, modified rectangular, scored tablets, debossed on one side with “A-007” and “5”.

Bottles of 30	NDC 59148-007-13
Blister of 100	NDC 59148-007-35

The 10-mg ABILIFY tablets are pink, modified rectangular tablets, debossed on one side with “A-008” and “10”.

Bottles of 30	NDC 59148-008-13
Blister of 100	NDC 59148-008-35

The 15-mg ABILIFY tablets are yellow, round tablets, debossed on one side with “A-009” and “15”.

Bottles of 30	NDC 59148-009-13
Blister of 100	NDC 59148-009-35

The 20-mg ABILIFY tablets are white, round tablets, debossed on one side with “A-010” and “20”.

Bottles of 30	NDC 59148-010-13
Blister of 100	NDC 59148-010-35

The 30-mg ABILIFY tablets are pink, round tablets, debossed on one side with “A-011” and “30”.

Bottles of 30	NDC 59148-011-13
Blister of 100	NDC 59148-011-35

U.S. Patent Nos. 4,734,416 and 5,006,528

Storage

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

Marketed by Otsuka America Pharmaceutical, Inc, Rockville, MD 20850 USA and Bristol-Myers Squibb Co, Princeton, NJ 08543 USA

Manufactured by Otsuka Pharmaceutical Co, Ltd, Tokyo, 101-8535 Japan

Distributed by Bristol-Myers Squibb Co, Princeton, NJ



XXXXXXXXXX XXXXXXXXXXXX

Issued XXXX 2002

© 2002, Otsuka Pharmaceutical Co, Ltd, Tokyo, 101-8535 Japan