

NASIVIN

Decongestant Nasal Drops

Composition:

NASIVIN for Paediatric use is a 0.025% aqueous solution of Oxymetazoline Hydrochloride.

NASIVIN for adults and school children is a 0.05% aqueous solution of Oxymetazoline Hydrochloride.

Properties:

When used locally, **NASIVIN** has an intensive vasoconstricting and decongesting effect on the nasal mucosa. The effect begins within a few minutes and continues for 6-8 hours.

Indications:

Acute colds. Paranasal Sinus inflammation. Syngitis.

Dosage and instructions for use:

NASIVIN 0.025% solution (Paediatric nasal drops) may only be taken by children over 1 year of age. With head bent, put forwards 1-2 drops of **NASIVIN** paediatric 0.025% from the dropper into each nostril 2-3 times daily or as directed by the physician.

NASIVIN 0.05% solution (adult nasal drops) may only be used by adults

and school children. Put into each nostril 1-2 drops of **NASIVIN** from the dropper 2-3 times daily.

Precautions:

Do not use **NASIVIN** more often than recommended, only use temporarily for up to 4 weeks if without a doctor's prescription. As with almost all drugs restrained use in the early stage of pregnancy.

Presentation:

10 mL bottle with dropper.

THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep all medicines out of the reach of children.



Produced by: **Julphar**
Gulf Pharmaceutical Industries,
Ras Al Khaimah, U. A. E.
Under licence from MERCK