

MULTI-VITAMIN CAPSULES, DROPS AND SYRUP

Vitamin deficiencies seldom occur singly and are most marked in growing children and adolescents, lactating mothers, those on weight loss, vegan, vegetarian, or poorly balanced diets, and during and after illnesses. During pregnancy there is often a requirement for additional vitamins but it is important to seek guidance from a doctor or antenatal clinic*.

CONCAVIT products provide the essential factors required in a dietary supplement to aid the prevention of vitamin deficiencies.

CONCAVIT DROPS are especially suitable for children and infants. They are highly concentrated to enable them to be mixed into drinks or blended with semi-solid and pureed infants' foods.

CONCAVIT CAPSULES

Each capsule contains:-

Vitamin A BP	5000.00	iu
Calciferol BP	500.00	iu
Ascorbic Acid BP	40.00	mg
Thiamine Mononitrate USP	2.50	mg
Riboflavine BP	2.50	mg
Pyridoxine Hydrochloride BP	1.00	mg
Nicotinamide BP	20.00	mg
Calcium Pantothenate USP	5.00	mg
Tocopheryl Acetate BP	2.00	iu

DOSAGE Adults and Children - 1 capsule daily

PACKING 25 and 100 capsules

PL 0400/5009

CONCAVIT SYRUP

Each 5ml contains:-

Vitamin A BP	5000.00	iu
Calciferol BP	500.00	iu
Ascorbic Acid BP	19.35	mg
Sodium Ascorbate USP	34.44	mg
Thiamine Hydrochloride USP	2.00	mg
Riboflavine BP	1.00	mg
(as Riboflavine Sodium Phosphate BP)	(1.37	mg)
Pyridoxine Hydrochloride BP	1.00	mg
Nicotinamide BP	12.50	mg
d-Panthenol	2.00	mg

DOSAGE Adults and Children - 5 ml per day

PACKING Bottles of 150ml,

PL 0400/5012

CONCAVIT DROPS

Each 0.5ml contains:-

Vitamin A BP	5000.00	iu
Calciferol BP	500.00	iu
Ascorbic Acid BP	19.35	mg
Sodium Ascorbate USP	34.44	mg
Thiamine Hydrochloride USP	2.00	mg
Riboflavine BP	1.00	mg
(as Riboflavine Sodium Phosphate BP)	(1.37	mg)
Pyridoxine Hydrochloride BP	1.00	mg
Nicotinamide BP	12.50	mg
d-Panthenol	2.00	mg

DOSAGE Adults and Children - 0.5ml (15 drops) per day

PACKING Bottles of 15ml, supplied with special dropper.

PL 0400/5010

Concavit is a medicine.

Do not exceed the stated dose.

Keep all medicines out of the reach of children.

Store between 4°C and 25°C

*If you are pregnant or likely to become pregnant you are advised to consult your Doctor or antenatal clinic before taking Vitamin A supplements. They will advise you on the best way to manage your diet and on what food combinations will be best for you. Generally during pregnancy it is best to avoid concurrent use of vitamin A supplements and liver.

Made in ENGLAND

WALLACE Manufacturing Chemists Ltd

52 Queen Anne Street, London W1M 9LA