

Anginova® Lozenges

This is a medicament
Medicament is a product which affects your health and its consumption contrary to instructions is dangerous for you. Follow strictly the doctor's prescription, the method of use and the instruction of the pharmacist who sold the medicament. – The doctor and the pharmacist are the experts in medicines, their benefits and risks. – Do not by yourself interrupt the period of treatment prescribed. – Do not repeat the same prescription without consulting your doctor. Keep all medicaments out of reach of children.



What is Anginova and when is it taken?

Anginova is a medicine used to treat inflammation and infections of the bucco-pharyngeal regions, especially for sore throat, pain on swallowing, thrush (Candida infection) and aphtous stomatitis, as well as for adjuvant therapy in angina. Each lozenge of Anginova contains two active substances. Dequalinium chloride is effective against most of the bacteria and fungi which cause infections of the mouth and throat as well as against yeasts (Candida). Lidocaine is analgetic and relieves pain on swallowing.

When must Anginova not be taken?

Anginova must not be taken by patients with known hypersensitivity towards one of the ingredients. Not appropriate for children under the age of 12.

When is caution advised during the intake of Anginova?

Since its efficacy is reduced by toothpaste, Anginova should not be taken shortly before or after brushing your teeth. Consult your doctor in case of high fever or if your symptoms do not improve within one week. Inform your doctor or pharmacist if you

- suffer from other illnesses,
- have allergies
- are using or taking other medicines (including non-prescription (over-the-counter) medicines)

Can Anginova be taken during pregnancy or whilst breast-feeding?

If you are pregnant or intend to become pregnant, you should avoid taking any medicine as a precautionary measure or consult your doctor or pharmacist. During pregnancy and lactation you should consult your doctor before taking Anginova.

How do you take Anginova?

Adults and young people from 12 years of age: Unless prescribed otherwise by the doctor, allow one lozenge to dissolve slowly in the mouth every 2 hours. The application and safety in children under the age of 12 has not yet been tested. The daily dose should not exceed 12 lozenges. Do not change the prescribed dosage. Please consult your doctor or pharmacist if you think that the medicine is too strong or too weak.

What are the possible side effects of Anginova?

The following side effects may occur when taking Anginova: Occasionally, hypersensitivity reactions have been observed. If you experience a hypersensitivity reaction, you should stop taking the medicine. Rarely, irritation of the oral mucosa has been observed. If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Other points to note

Keep the medicine out of the reach of children. Do not store above 30 °C. The medicine can only be taken up to the expiry date “EXP” indicated on the package. Please consult your doctor or pharmacist for more detailed information.

What does Anginova contain?

1 g lozenge contains:
Active substances: Dequalinium chloride 0,25 mg, Lidocaine hydrochloride 1 mg
Excipients: Lactose anhydrous, Sodium cyclamate, Peppermint aromas, Povidone, Glycerol dibehenate

Where can you obtain Anginova?

What pack sizes are available?
Anginova is available from pharmacies without prescription.
Pack sizes of 20 lozenges.

Manufactured by:

Rottendorf Pharma GmbH, Ostenfelder Strasse 51-61, 59320 Ennigerloh, Germany

For:



Medinova AG, Eggbühlstr. 28, 8050 Zurich, Switzerland

