

OVERDOSE

The symptoms of adverse reaction may occurred with overdose of Microclismi.

In case of overdose, the support therapy for relief of adverse reaction should be instituted. Promptly carry out the measures that can prevent the absorption of drug and rapidly eliminate the drug out of the body.

STORAGE

Store at temperature not exceeding 25°C, protected from sunlight.

EXPIRY

60 months from the manufacturing date.

The expiry date shown refers to the unopened packet, stored correctly.

HOW SUPPLIED

6 bottles/box.

SPECIFICATION: Manufacturer.

WARNINGS

Read carefully the leaflet before use

For further information, please contact your doctor

Keep out of the reach of children.

Manufacturers: ZETA FARMACEUTICI S.p.A.

Via Galvani,10 - 36066 - Sandrigo (VI) - Italy

Marketing Authorization Holder: Industria Farmaceutica NOVA ARGENTIA S.p.A.

Via Lovanio, 5 - 20121 Milano - Italy.

MICROCLISMI GLYCEROL, CHAMOMILE AND MALLOW

9 g and 3 g micro-enemas

COMPOSITION

Each children micro-enema contains:

Active ingredient: Glycerol 2,25 g, Chamomile Fluid Extract 0,10 g, Mallow Fluid Extract 0,10 g

Inactive ingredient: Wheat Starch 0,02 g, Purified Water q.s.

Each adults micro-enema contains: Glycerol 6,75 g, Chamomile Fluid Extract 0,30 g, Mallow Fluid Extract 0,30 g

Inactive ingredient: Wheat Starch 0,06 g, Purified Water q.s.

INDICATION

Short-term treatment of constipation in order to facilitate and to quick intestinal evacuation.

DOSAGE AND ADMINISTRATION

Rectal administration.

1-2 micro-enemas in 24 hours.

Don't use more than two micro-enemas per administration and don't exceed recommended dose.

Dosage may be adjusted according to age and symptom.

CONTRAINDICATION

Hypersensitivity to active ingredients or excipients.

Laxatives are contraindicated in patients with acute or of uncertain origin abdominal pain, nausea, vomiting gastrointestinal obstruction or stenosis.

Patients with rectal bleeding of unknown origin, advanced state of dehydration.

The use of enemas should be avoided in patients suffering from acute painful haemorrhoids with bleeding.

ADVERSE REACTION

Adverse effects are primarily due to dehydrating action of glycerol. When given rectally may cause irritation.

Inform your doctor with side effects when using medicine.

GENERAL PRECAUTION

Consult a doctor if an obstinate constipation persist.

DRUG INTERACTIONS

No known interactions with other medicines.

USE ON PREGNANCY OR LACTATION

No data are available.

F05770103001