

LOMEXIN®

Cream, spray, sprinkling powder,
dermo-foam

Fenticonazole nitrate
Dermatological preparations

COMPOSITION

- **Cream** 100 g contain:

Fenticonazole nitrate 2g.

Excipients: propylene glycol, hydrogenated lanolin, almond oil, polyglycolic ester of fatty acids, cetyl alcohol, glyceryl monostearate, sodium edetate, purified water.

- **Spray** 100 g contain:

Fenticonazole nitrate 2g.

Excipients: alcohol, propylene glycol, purified water.

- **Sprinkling powder** 100 g contain:

Fenticonazole nitrate 2g.

Excipients: precipitated silica, zinc oxide, talc.

- **Dermo-foam** 100 ml of solution contain:

Fenticonazole nitrate 2g.

Excipients: propylene glycol, alcohol, sodium lauryl ether sulfate solution, alkylamidobetaine solution, purified water.

PRESENTATIONS

2% cream: tube of 30 g.

2% spray: bottle of 30 ml.

2% sprinkling powder: bottle of 50g.

2% dermo-foam: bottle of 100 ml with dispenser.

PROPERTIES

Lomexin exerts fungistatic and fungicidal action on dermatophytes (responsible for most mycotic skin infections), pathogenic yeasts, dimorphous fungi and moulds. In addition, Lomexin is active against gram positive bacteria and is therefore indicated in mycoses associated with bacterial superinfections.

INDICATIONS

- Dermatomycosis caused by dermatophytes (*Trichophyton*, *Microsporum*, *Epidermophyton*) at different locations: tinea capitis, tinea corporis, tinea cruris, tinea pedis (athlete's foot), tinea manuum, tinea faciei, tinea barbae, tinea unguium.

- Cutaneous candidiasis (intertrigo, perleche, face candidiasis, "napkin" rash, perineal and scrotal candidiasis); balanitis, balanoposthitis, onychia and paronychia.

- *Pytirisias versicolor* (caused by *Pytirosporum orbiculare* and *P. ovale*).

- Otomycosis (due to *Candida* or moulds) only if no lesions of eardrum are present.

- Erythrasma.

- Mycosis with bacterial superinfections (caused by Gram-positive bacteria).

CONTRAINDICATIONS

Ascertained hypersensitivity to the product.

Although transcutaneous and vaginal absorption of fenticonazole is very poor, use of the preparation is not recommended in pregnancy.

WARNINGS

The use, especially if on a long-term basis, of products for topical application, could induce sensitization phenomena. In this case, discontinue the treatment and seek the doctor's advice. Do not expose to sun rays or UVA treatment after the dermo-foam application.

USE AND DOSAGE

Lomexin should be applied once or twice daily, according to medical instructions, on the washed and dried injured area using the suitable formulation according to the site of lesion.

- **Cream:** Indicated for the treatment of glabrous skin, cutaneous folds and mucous membranes: to apply, gently rub the cream on the injured area.

Especially indicated for dry mycosis: *Pytirisias versicolor*, erythrasma, onychomycosis (in onychomycosis the cream should be applied by occlusive dressing); the cream is also suitable in male genital mycosis.

- **Spray:** Is indicated in the forms affecting the scalp and for any skin areas covered with hair. This is a practical and easy-to-use formulation for the treatment of large mycosis and areas which are not easily reached.

- **Sprinkling powder:** It is particularly indicated for athlete's foot and in general for intertriginous areas and moist lesions, either alone or as adjuvant to the cream treatment. In case of foot infections, it is recommended to sprinkle Lomexin powder inside both socks and shoes.

In the case of tinea pedis and onychomycosis, the treatment should be prolonged for 1 to 2 weeks after healing of lesions in order to avoid re-infections.

- **Dermo-foam:** It is particularly indicated in the treatment of *Pytirisias versicolor* and of wide spread dermatomycosis of hairless skin; for *Pytirisias versicolor*, apply directly on wet skin by means of a sponge in the evening, allow to dry overnight and wash away on the morning after. Due to its physiological pH, Lomexin dermo-foam is a particularly recommended cleanser during the treatment of infections, to prevent further skin irritation. Lomexin is not greasy, does not soil and can be easily removed with water.

SIDE EFFECTS

Lomexin (fenticonazole) is generally well tolerated by the skin and mucous membranes: only exceptionally mild and transient erythematous reactions have been reported. After topical application a slight burning sensation may occur, usually subsiding soon. Should more persistent irritation occur or resistant microorganisms develop, suspend the treatment and seek the doctor's advice. Due to Lomexin poor absorption, no systemic effects should develop, provided the above instructions are carefully observed.

STORAGE CONDITIONS

Spray: Store in a dry place below 30°C, protected from light.

Other forms: Store in a dry place below 25°C, protected from light.

Do not refrigerate.

Do not use after expiry date. The expiry date refers to the last day of that month.

Not all presentations may be marketed.

Keep medicament out of reach and sight of children.

This is a medicament

- A Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.

- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.

- The doctor and the pharmacist are experts in medicine, its benefits and risks.

- Do not by yourself interrupt the period of treatment prescribed.

- Do not repeat the same prescription without consulting your doctor.

Manufactured by
ALGORITHM S.A.L. Zouk Mosbeh
Lebanon
Under License from Recordati

 RECORDATI

® Registered Trademark.