

LOMEXIN® T

1000 mg ovule
Fenticonazole nitrate

COMPOSITION

Each ovule contains:

Active ingredient:

Fenticonazole nitrate 1000 mg.

Excipients:

Liquid paraffin, white soft paraffin, soya lecithin, gelatin, glycerin, titanium dioxide, sodium ethylhydroxybenzoate, sodium propylhydroxybenzoate.

PRESENTATION

Pack of one 1000 mg ovule.

PROPERTIES

Lomexin® exerts fungistatic and fungicidal actions on dermatophytes (responsible for most mycotic skin infections), pathogenic yeasts, dimorphous fungi and moulds. In addition Lomexin® is active against gram-positive bacteria and is therefore indicated in mycoses associated with bacterial superinfections.

Lomexin® is also active, both in vitro and vivo, against trichomonas vaginalis.

INDICATIONS

Trichomonas vaginalis.

Vaginal infections sustained by mixed forms of trichomonas vaginalis and candida albicans.

Genital candidiasis (vulvovaginitis, colpitis, infectious fluor).

CONTRAINDICATION

Ascertained hypersensitivity to the product.

WARNINGS

Although transcutaneous and vaginal absorption of Fenticonazole is very poor, use of the preparation is not recommended in pregnancy.

DOSAGE AND ADMINISTRATION

Trichomonas or mixed (trichomonas and candida albicans) vaginal infections:

One 1000 mg ovule followed by a second administration 24 hours later, when necessary.

Candida albicans infections:

One single 1000 mg ovule administration in the evening. Should the symptoms persist, a second administration may be repeated after three days.

The ovules must be introduced deep into the vagina and pushed well up to the fornix.

To avoid re-infection, it is recommended that the partner undergoes concurrent local (glans and prepuce) treatment with Lomexin® cream.

SIDE EFFECTS

Lomexin® (Fenticonazole) is generally well tolerated by the mucous membranes; only exceptionally mild and transient erythematous reactions have been reported. After topical application or intravaginal administration, a slight burning sensation may occur, usually subsiding soon. Should more persistent irritation occur or resistant micro-organism develop, suspend the treatment and seek the doctor's advice. Due to Lomexin® poor absorption, no systemic effects should occur, provided the above instructions are carefully observed.

Do not use after expiry date.

STORAGE CONDITIONS

Store in a dry place below 25°C, protected from light.

Do not refrigerate.

This is a medicament

-A medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.

-Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.

-The doctor and the pharmacist are experts in medicine, its benefits and risks.

-Do not by yourself interrupt the period of treatment prescribed.


-Do not repeat the same prescription without consulting your doctor.

KEEP MEDICAMENT OUT OF CHILDREN'S REACH

Packed in Zouk mosbeh Lebanon, by

ALGORITHM S.A.L.

Under License from Recordati

 Registered Trademark.