

Pediatric Drops

Restropinal[®] is specifically formulated for the symptomatic relief of gastro-intestinal spasm in infants and children.

COMPOSITION

Each cc (20 drops) contains:

Methscopolamine nitrate 0.1 mg

ACTION

Antispasmodic: Methscopolamine nitrate has a powerful and rapid effect. It is more efficient and acts more quickly than Atropine or Methylatropine, with the following advantages:

- Marked anticholinergic activity.
- Absence of the central effect of Scopolamine as this drug does not penetrate the blood-brain barrier.
- More selective antispasmodic effect on the smooth muscle.
- No toxic effects or tachycardia at usual doses.

Sedative: Butabarbital (acid) is a barbiturate of medium duration of action. A small dose provides a mild sedative effect to the stressed child.

INDICATIONS

Spastic conditions of the gastro-intestinal tract such as intestinal colic, pylorospasm, hypertrophic pyloric stenosis.

Functional nausea and vomiting.

DOSAGE

Before each feeding five to six times daily.

Birth to one year: Two drops per kilogram of body weight 5 to 6 times daily.

One to two years: Half teaspoonful 5 to 6 times daily. Two to six years: One teaspoonful 5 to 6 times daily.

CONTRAINDICATIONS

Glaucoma and cardiac arrhythmias.

SIDE EFFECTS

When the recommended dose is exceeded, and in some patients with average therapeutic doses, one or more of the following side effects may occur and require reduction of the dose: dryness of the mouth, blurring of vision, constipation, difficulty in starting urination, weakness, headache, drowsiness and slow pulse rate. Should any troublesome or unusual symptoms develop, a doctor should be consulted.

CAUTION

Restropinal[®] Pediatric Drops should not be used in case of organic obstruction of the gastro-intestinal tract.

Do not exceed the prescribed dose.

HOW SUPPLIED

Restropinal[®] Pediatric Drops is supplied in 25 cc. bottles with calibrated dropper.

20 drops from the calibrated dropper measure 1 cc.

STORAGE CONDITIONS

Store in a dry place below 30°C protected from light. Do not refrigerate.

Do not use after expiry date.

To be dispensed only by or on the prescription of a physician.

This is a medicament

- -A medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- -Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- -The doctor and the pharmacist are experts in medicine, its benefits and risks.
- -Do not by yourself interrupt the period of treatment prescribed.
- -Do no repeat the same prescription without consulting your doctor.

KEEP MEDICAMENT OUT OF REACH OF CHILDREN