

CLOREN

Diclofenac

FORM AND PRESENTATION:

Suppository for Adults of 100 mg: Box of 8. Nonsteroidal Anti-inflammatory Drug (NSAID)

INDICATIONS:

- *Long-term symptomatic treatment:*

Chronic inflammatory forms of rheumatism, rheumatoid polyarthritis and ankylosing spondylarthritis

- *Short-term symptomatic treatment:*

Acute attacks of tendinitis, bursitis, microcrystalline arthritis, arthrosis, lumbago and painful syndromes of the vertebral column.

ADMINISTRATION:

1 suppository/day at night before bedtime.

TERM OF TREATMENT:

Do not exceed a period of 2 to 3 days.

CONTRAINDICATIONS:

- Antecedent of allergies or asthma attacks caused by intake of diclofenac or any other substance of similar effects, such as other NSAID, aspirin.
- Hypersensitivity to one of the excipients.
- Developing gastroduodenal ulcer.
- Severe hepatocellular deficiency.
- Severe kidney deficiency.
- Trouble of hemostasis or during anticoagulant treatment.
- Antecedent of rectum inflammation.
- Pregnancy (3rd trimester).

INTERACTIONS

The simultaneous administration of Diclofenac with the products listed below necessitates a strict surveillance of the clinical and biological condition of the patient.

Inadvisable combinations:

- Oral anticoagulant
- Other NSAID (including high doses of salicylates)
- Heparines (by parenteral route)

- Lithium
- Methotrexate (at doses > 15 mg / week)
- Ticlopidine

Combinations necessitating precautions:

- Diuretics, inhibitors of the conversion enzyme and angiotensine II.
- Methotrexate (at doses < 15 mg / week)
- Pentoxifylline
- Zidovudine

Combinations to be taken into consideration:

- Betablocking agents
- Cyclosporine
- Intrauterine device
- Thrombolytic agents

ADVERSE EFFECTS:

- Infrequent gastrointestinal upsets at the start of treatment: abdominal pain, nausea, diarrhea, dyspepsia and anorexia.
- Hypersensitive reactions mainly dermatological reactions: skin rash, urticaria and eczema.

PRECAUTIONS:

This medicine ought to be taken under medical surveillance.

PREGNANCY AND LACTATION:

The prescription of NSAID is not recommended unless necessary during the first five months of pregnancy. Starting the 6th month and on, this medicine is contraindicated. NSAID passes into the breast milk, it is better to avoid the use of such substances during lactation.

CONSERVATION:

All medicines should be kept out of children's reach.

Do not use when expired, expiry date on the box.

Keep in a dry cool place at room temperature under 30°C.



CHAPHA (Chalhoub Pharmaceuticals) S.A.L.

Manufactured in Lebanon