

INDOLOR[®]

DESCRIPTION

INDOLOR is an oral rinse containing 0.2% Chlorhexidine digluconate + 25% Alcohol.

INDICATIONS AND USAGE

Chlorhexidine is an antiseptic & disinfectant against a wide range of bacteria, some fungi & some viruses. It is used in the treatment & prevention of gingivitis, prevention of plaque & in the maintenance of oral hygiene.

- **INDOLOR** Oral Rinse is indicated for use between dental visits as part of a professional program for the treatment of gingivitis as characterized by redness and swelling of the gingivae, including gingival bleeding upon probing.
- **INDOLOR** is useful in the management of recurrent oral ulceration & oral candidal infections
- **INDOLOR** may also be used to promote gingival healing after periodontal

CONTRAINDICATIONS

INDOLOR Oral Rinse should not be used by persons who are known to be hypersensitive to chlorhexidine digluconate

ADVERSE REACTIONS

The most common side effects associated with chlorhexidine digluconate oral rinses are :

- an increase in staining of teeth & other oral surfaces
- an increase in calculus formation
- an alteration in taste perception
- Discoloration of the teeth & tongue may occur. This discoloration disappears after treatment is discontinued.

Minor irritation and superficial desquamation of the oral mucosa have been noted in patients using chlorhexidine digluconate oral rinses, particularly among children.

DOSAGE AND ADMINISTRATION

INDOLOR is not intended for ingestion & should be expectorated after rinsing.

Place one table spoonful of **INDOLOR** in half a cup of water and rinse the mouth for about 1 minute.

STORAGE

Store below 25° C

Protect from light

Do not swallow this product

COMPOSITION

Each 100 ml contains: Chlorhexidine digluconate 0.2g

Inactive Ingredients: Alcohol - Glycerine

Saccharine-Propylene Glycol-Mint Flavor-Dye Blue

Pack

100 ml Bottle

THIS IS A MEDICAMENT

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you .
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep all medicines out of the reach of children

(ان هذا الدواء)

- الدواء مستحضر يؤثر على صحتك واستهلاكه خلافا للتعليمات يعرضك للخطر .
- اتبع بدقة وصفة الطبيب وطريق الاستعمال المنصوص عليها وتعليمات الصيدلاني الذي صرفها لذلك.
- فالطبيب والصيدلاني هما الخبيران بالدواء وبنفعه وضرره
- لا تقطع مدة العلاج المحددة لك من تلقاء نفسك
- لا تكرر صرف الدواء بدون وصفة طبية

أحفظ الدواء بعيداً عن متناول الأطفال

PHARMADEX S.A.L.
KAHALE - LEBANON



Revised May 2006