

PHARMALINE

Decalyn

Bactericidal, Fungistatic Mouth Antiseptic & Anti-inflammatory

DESCRIPTION

Decalyn is a combination product of Dequalinium chloride and Lidocaine HCL formulated into pleasant tasting sugar-free lozenges.

ACTIONS

Dequalinium chloride is a wide spectrum antiseptic with bactericidal and fungistatic properties. It has a rapid effect on organisms causing mixed infections of the mouth and pharynx. Even bacterial strains which are resistant to antibiotics are eliminated. The added Lidocaine HCL component produces a mild local anesthesia, which relieves the painful symptoms accompanying mouth and throat infection and inflammation.

INDICATIONS

Local treatment of painful infections of the mouth and throat including anginous infections, pharyngitis, laryngitis, and candidiasis of the buccopharyngeal cavity. It is also indicated in the treatment of inflammations of the mucous membrane and gums including gingivitis, ulcerative stomatitis, and aphtous stomatitis.

Decalyn can also be used for post-operative applications following tonsillectomy and dental extractions, to promote smooth healing. In addition, *Decalyn* can be administered as prophylaxis in times of increased infections.

CONTRAINDICATIONS

Decalyn is contraindicated in patients with any known hypersensitivity to any of the two active ingredients or to local anesthetics.

Decalyn is also contraindicated in children below 6 years of age.

PRECAUTIONS

Decalyn should not be used for long periods of time. If the symptoms persist for longer than 5 days, the medical condition should be re-evaluated.

ADVERSE REACTIONS

Possible temporary tingling in the tongue.

DOSAGE & ADMINISTRATION

1 lozenge to be dissolved slowly in the mouth every 2 to 3 hours.

DRUG INTERACTIONS

None known.

AVAILABILITY

Boxes of 20 lozenges, each containing Dequalinium Chloride 0,25 mg, Lidocaine HCl 1 mg, Excipient q.s. 1 lozenge.

Reg Lebanon 973864



* Decalyn is a Trade Mark.