

# Spasfon<sup>®</sup>

phloroglucinol

## IDENTIFICATION OF MEDICINE

### Trade name of the medicinal product :

SPASFON, suppository.

### Qualitative and quantitative composition :

Hydrated phloroglucinol ..... 150,00 mg  
corresponding amount  
of anhydrous phloroglucinol ..... 116,70 mg  
Trimethylphloroglucinol ..... 150,00 mg  
Excipients : hard fat per suppository.

### Pharmaceutical form :

Suppository - pack of 10.

### Pharmaco-therapeutic classification :

MUSCULOTROPIC ANTISPASMODIC

(A : digestive system and metabolism)

(G : genitourinary system)

### Name and address of distributor and manufacturer :

Laboratoire L. LAFON - 19, avenue du Professeur Cadiot  
B.P. 22 - 94701 MAISONS-ALFORT (FRANCE).

## WHEN SHOULD THIS MEDICINE BE USED

### (Therapeutic indications)

This medicine is intended to be used for the treatment of  
spasmodic pain arising from the intestine, biliary tract,  
bladder and uterus.

## BEWARE !

### When should this medicine not be used :

#### (Contraindications)

This medicine MUST NOT BE TAKEN in case of allergy to  
any of its constituents.

IF IN DOUBT, IT IS ESSENTIAL TO SEEK THE ADVICE OF  
YOUR DOCTOR OR YOUR PHARMACIST.

### Precautions :

IF IN DOUBT, DO NOT HESITATE TO SEEK THE ADVICE OF  
YOUR DOCTOR OR YOUR PHARMACIST.

### Drug interactions and other forms of interaction :

IN ORDER TO AVOID POSSIBLE INTERACTIONS BETWEEN  
SEVERAL MEDICINES, YOU SHOULD ALWAYS INFORM YOUR  
DOCTOR OR PHARMACIST OF ANY ANOTHER TREATMENT  
THAT YOU MAY BE TAKING.

### Pregnancy - Breast feeding :

This medicine should be taken during pregnancy only  
if necessary. Women should not take this medicine during  
breast feeding.

AS A MATTER OF GENERAL PRINCIPLE, YOU SHOULD  
ALWAYS SEEK THE ADVICE OF YOUR DOCTOR OR  
PHARMACIST BEFORE TAKING ANY MEDICINE IF YOU ARE  
PREGNANT OR BREAST FEEDING.

## HOW TO USE THIS MEDICINE

### Dosage :

3 suppositories a day.

### Method and route of administration :

By rectum.

## UNDESIRABLE AND TROUBLESOME EFFECTS

### (Adverse effects)

IN COMMON WITH ALL ACTIVE SUBSTANCES, THIS MEDICINE  
MAY CAUSE TROUBLESOME EFFECTS OF VARYING DEGREES IN  
CERTAIN INDIVIDUALS :

- a very few allergic skin reactions have been reported.  
TELL YOUR DOCTOR OR PHARMACIST ABOUT ANY OTHER  
UNDESIRABLE AND TROUBLESOME EFFECT NOT MENTIONED  
IN THIS PACKAGE INSERT INFORMATION.

## STORAGE

**Do not exceed the expiry date shown on the outer  
packaging.**

### Special storage precautions :

This medicine must be stored away from heat.

## DATE OF REVIEW OF PACKAGE INSERT INFORMATION

October 1996.